

Simplicity of a vertex operator algebra whose Griess algebra is the Jordan algebra of symmetric matrices

Hidekazu Niibori

Graduate School of Pure and Applied Sciences, University of Tsukuba,
Tsukuba, Ibaraki 305-8571, Japan (e-mail: niibori@math.tsukuba.ac.jp)

and

Daisuke Sagaki

Institute of Mathematics, University of Tsukuba,
Tsukuba, Ibaraki 305-8571, Japan (e-mail: sagaki@math.tsukuba.ac.jp)

Abstract

Let $r \in \mathbb{C}$ be a complex number, and $d \in \mathbb{Z}_{\geq 2}$ a positive integer greater than or equal to 2. Ashihara and Miyamoto introduced a vertex operator algebra $V_{\mathcal{J}}$ of central charge dr , whose Griess algebra is isomorphic to the simple Jordan algebra of symmetric matrices of size d . In this paper, we prove that the vertex operator algebra $V_{\mathcal{J}}$ is simple if and only if r is not an integer. Further, in the case that r is an integer (i.e., $V_{\mathcal{J}}$ is not simple), we give a generator system of the maximal proper ideal I_r of the VOA $V_{\mathcal{J}}$ explicitly.

1 Introduction.

Let $V = \bigoplus_{n \in \mathbb{Z}_{\geq 0}} V_n$ be a vertex operator algebra (VOA for short) over the field $\mathbb{C}$ of complex numbers with $Y(\cdot, z) : V \rightarrow (\text{End } V)[[z, z^{-1}]]$ the vertex operator. As usual, for each $v \in V$, we define $v_m \in \text{End } V$, $m \in \mathbb{Z}$, by: $Y(v, z) = \sum_{m \in \mathbb{Z}} v_m z^{-m-1}$. It follows from the axiom of a VOA that V_2 becomes a $\mathbb{C}$ -algebra with the product given by $u \cdot v := u_1 v \in V_2$ for $u, v \in V_2$. In addition, we know (see [FLM, §10.3] and also [M, §5]) that if $\dim V_0 = 1$ and $\dim V_1 = 0$, then the $\mathbb{C}$ -algebra V_2 is commutative (but not necessarily associative), which we call the Griess algebra of V .

Various kinds of commutative $\mathbb{C}$ -algebras appear as the Griess algebras of VOAs. The Griess algebra of the moonshine VOA $V^\natural$ is isomorphic to the 196884-dimensional, commutative $\mathbb{C}$ -algebra introduced by Griess [G], whose automorphism group is isomorphic to the Monster sporadic simple group (see [FLM, p.319]); the name ‘‘Griess algebra’’ is derived from this fact. Also, for a given associative, commutative $\mathbb{C}$ -algebra A equipped

with an A -invariant bilinear form, Lam [L1] constructed a VOA whose Griess algebra is isomorphic to A . Further, in [L2], [AM], [As], they constructed some VOAs whose Griess algebras are (simple) Jordan algebras; for the definition of a Jordan algebra, see §2.2 below. In this paper, we will mainly treat VOAs introduced in [L2, §4.1] and [AM], whose Griess algebras are isomorphic to the simple Jordan algebra $\text{Sym}_d(\mathbb{C})$ of symmetric matrices of size $d \in \mathbb{Z}_{\geq 2}$ with entries in $\mathbb{C}$. Because the Jordan algebra $\text{Sym}_d(\mathbb{C})$ has a strong connection to symmetric cones and zeta functional equations (see [FK]), Ashihara, Miyamoto (see [AM, Introduction]), and the authors of this paper expect that the results in [AM] and this paper contribute a VOA theoretical approach to the theory of symmetric cones and zeta functional equations.

An essential difference between the VOA introduced in [L2, §4.1] and the one introduced in [AM] (which we denote by $V_{\mathcal{J}}$) is their central charges. The central charge of the former is equal to the (fixed) positive integer $d \in \mathbb{Z}_{\geq 2}$, the size of symmetric matrices. On the other hand, the central charge of the later is equal to dr , where r is an arbitrary complex number (see Theorem 2.5 below). In general, the structure and the representation theory of a VOA deeply depends on its central charge. For example, it is well-known that the simplicity of a Virasoro VOA $M_{c,0}/\langle L_{-1}1 \rangle$ (with notation in [W]) and the rationality of a simple Virasoro VOA V_c (with notation in [W]) depend on their central charges c (see [W] and also [DMZ]). Moreover, rational Virasoro VOAs (i.e., V_c of special central charge c , such as $V_{1/2}$) and their irreducible modules play very important roles in the theory of VOAs. So, as in the case of Virasoro VOAs, it is quite natural and important to study how the VOA $V_{\mathcal{J}}$ introduced in [AM] depends on its central charge dr . In this paper, we study the condition of $r \in \mathbb{C}$ for the VOA $V_{\mathcal{J}}$ to be simple. The main result of this paper is the following theorem, which means that the simplicity of $V_{\mathcal{J}}$ also depends on its central charge.

Theorem. *Keep the notation above. The VOA $V_{\mathcal{J}}$ is simple if and only if r is not an integer.*

Many important VOAs are obtained as the nontrivial simple quotients of nonsimple VOAs (e.g., rational Virasoro VOAs, and VOAs associated to integrable highest weight modules over affine Lie algebras). So we are interested in the simple quotient $V_{\mathcal{J}}/I_r$ with $r \in \mathbb{Z}$ rather than the VOA $V_{\mathcal{J}}$. When we study the structure of $V_{\mathcal{J}}/I_r$ and its representation theory, it is very important to determine some relations in $V_{\mathcal{J}}/I_r$ induced by the maximal proper ideal I_r . In this paper, as a first step for studying the simple VOA $V_{\mathcal{J}}/I_r$, we will give a generator system of the maximal proper ideal I_r of the VOA $V_{\mathcal{J}}$ explicitly, whose elements are singular vectors for a certain Lie algebra $\mathcal{L}_r^{(1)}$, and have a high symmetry (see §6 below).

This paper is organized as follows. In §2, we recall the definition of the Griess algebra of a VOA, the definition of a Jordan algebra, and the construction of the VOA $V_{\mathcal{J}}$ introduced by Ashihara and Miyamoto [AM]. Then we state our main theorem (Theorem 2.6), and the plan how we prove the theorem. In §3–§6, following the plan, we will prove some key propositions (Propositions 3.1, 3.4, 4.1, 5.1, 6.1); our main theorem follows immediately from these propositions. In §6, we also give a generator system of the maximal ideal I_r of the VOA $V_{\mathcal{J}}$ explicitly when r is an integer, that is, when $V_{\mathcal{J}}$ is not simple.

Acknowledgments. The authors would like to express their sincere gratitude to Professor Masahiko Miyamoto, Professor Toshiyuki Abe, Professor Hiroki Shimakura, Professor Hiroshi Yamauchi, and Dr. Takahiro Ashihara for valuable discussions.

2 Vertex operator algebra whose Griess algebra is a Jordan algebra.

2.1 Griess algebras. Let $(V = \bigoplus_{n \in \mathbb{Z}_{\geq 0}} V_n, Y(\cdot, z), \mathbf{1}, \omega)$ be a vertex operator algebra (VOA for short), with $Y(\cdot, z) : V \rightarrow (\text{End } V)[[z, z^{-1}]]$ the vertex operator, $\mathbf{1} \in V_0$ the vacuum element, and $\omega \in V_2$ the Virasoro element (for the details about VOAs, see, e.g., [LL]). As usual, for each $v \in V$, we define $v_m \in \text{End } V$, $m \in \mathbb{Z}$, by: $Y(v, z) = \sum_{m \in \mathbb{Z}} v_m z^{-m-1}$. For $a, b \in V_2$, we define $a \cdot b := a_1 b$. Then it follows from the axiom of a VOA that $a \cdot b \in V_2$ for every $a, b \in V_2$, i.e., V_2 becomes a $\mathbb{C}$ -algebra with $\cdot$ the product. In addition, if $V_0 = \mathbb{C}\mathbf{1}$ and $V_1 = \{0\}$, then the $\mathbb{C}$ -algebra V_2 is commutative (see [FLM, §10.3] and also [M, §5]). In this case, we call V_2 the Griess algebra of V . Note that the Griess algebra of a VOA is not necessarily associative.

2.2 Jordan algebras. Let us recall the definition of a Jordan algebra. For the details about Jordan algebras, see, e.g., [Al1], [Al2], and [J].

Definition 2.1. Let J be a $\mathbb{C}$ -algebra with the product $a \cdot b$ ($a, b \in J$). The $\mathbb{C}$ -algebra J is called a Jordan algebra if $a \cdot b = b \cdot a$ and $a^2 \cdot (b \cdot a) = (a^2 \cdot b) \cdot a$ hold for every $a, b \in J$.

Let $\text{Sym}_d(\mathbb{C})$ be the set of symmetric matrices of size $d \in \mathbb{Z}_{\geq 2}$ with entries in $\mathbb{C}$. It is well-known that $\text{Sym}_d(\mathbb{C})$ becomes a (simple) Jordan algebra, where the product is given by: $A \cdot B = \frac{1}{2}(AB + BA)$ for $A, B \in \text{Sym}_d(\mathbb{C})$ (see also [L2, Theorem 6 B]).

2.3 VOA $V_{\mathcal{J}}$. Let (and fix) $d \in \mathbb{Z}_{\geq 2}$. In this subsection, we recall a VOA $V_{\mathcal{J}}$ introduced by Ashihara and Miyamoto [AM], whose Griess algebra is isomorphic to the Jordan algebra $\text{Sym}_d(\mathbb{C})$ of symmetric matrices.

Let $\widehat{\mathfrak{h}}$ be an (infinite-dimensional) vector space over $\mathbb{C}$ with a linear basis $\{v^i(m) \mid 1 \leq i \leq d, m \in \mathbb{Z}\} \cup \{\mathbf{c}\}$, and define a Lie bracket on $\widehat{\mathfrak{h}}$ by:

$$\begin{aligned} [v^i(m), v^j(n)] &= \delta_{m+n,0} \delta_{i,j} m\mathbf{c} \quad \text{for } 1 \leq i, j \leq d \text{ and } m, n \in \mathbb{Z}, \\ [\mathbf{c}, \widehat{\mathfrak{h}}] &= \{0\}. \end{aligned} \tag{2.1}$$

Denote by $U(\widehat{\mathfrak{h}})$ the universal enveloping algebra of the Lie algebra $\widehat{\mathfrak{h}}$, and let $U(\widehat{\mathfrak{h}})/\langle \mathbf{c} - 1 \rangle$ be the quotient algebra with respect to the two-sided ideal $\langle \mathbf{c} - 1 \rangle$ of $U(\widehat{\mathfrak{h}})$ generated by $\mathbf{c} - 1 \in U(\widehat{\mathfrak{h}})$. We define a subspace $\mathcal{L}$ of $U(\widehat{\mathfrak{h}})/\langle \mathbf{c} - 1 \rangle$ as follows: For $1 \leq i, j \leq d$ and $m, n \in \mathbb{Z}$, we set

$$v^{ij}(m, n) := v^i(m)v^j(n) \mod \langle \mathbf{c} - 1 \rangle. \tag{2.2}$$

Let

$$\begin{aligned} \mathcal{B} := \{ &v^{ii}(m, n) \mid 1 \leq i \leq d \text{ and } m, n \in \mathbb{Z} \text{ with } m \leq n \} \cup \\ &\{v^{ij}(m, n) \mid 1 \leq i < j \leq d \text{ and } m, n \in \mathbb{Z}\}. \end{aligned}$$

Then it follows from the Poincaré-Birkhoff-Witt theorem that $\mathcal{B} \cup \{1 \in U(\widehat{\mathfrak{h}})/\langle \mathbf{c} - 1 \rangle\}$ is a linearly independent subset of $U(\widehat{\mathfrak{h}})/\langle \mathbf{c} - 1 \rangle$. We set

$$\mathcal{L} := (\text{Span}_{\mathbb{C}} \mathcal{B}) \oplus \mathbb{C} \subset U(\widehat{\mathfrak{h}})/\langle \mathbf{c} - 1 \rangle.$$

Remark 2.2. Let $1 \leq i, j \leq d$, and $m, n \in \mathbb{Z}$. It can be easily seen from the definition (2.1) of the Lie bracket on $\widehat{\mathfrak{h}}$ that

$$\begin{aligned} v^{ij}(m, n) &= v^{ji}(n, m) \quad \text{if } i \neq j, \text{ or if } i = j \text{ and } m \neq -n, \\ v^{ii}(m, -m) &= v^{ii}(-m, m) + m. \end{aligned} \tag{2.3}$$

In particular, $v^{ij}(m, n) \in \mathcal{L}$ for all $1 \leq i, j \leq d$ and $m, n \in \mathbb{Z}$.

We see by direct computation that $[x, y] = xy - yx$ is contained in $\mathcal{L}$ for every $x, y \in \mathcal{L}$, and hence $\mathcal{L}$ becomes a Lie algebra with respect to the natural Lie bracket. Now, let (and fix) $r \in \mathbb{C}$ be an (arbitrary) complex number. For each $x, y \in \mathcal{L}$, we define

$$[x, y]_r := \pi_1([x, y]) + r\pi_2([x, y]), \tag{2.4}$$

where $\pi_1 : \mathcal{L} \rightarrow \text{Span}_{\mathbb{C}} \mathcal{B}$ and $\pi_2 : \mathcal{L} \rightarrow \mathbb{C}$ denote projections from $\mathcal{L}$ onto $\text{Span}_{\mathbb{C}} \mathcal{B}$ and $\mathbb{C}$, respectively. Then we know from [AM, §2.1] that $[\cdot, \cdot]_r$ is a Lie bracket on $\mathcal{L}$. Let us denote by $\mathcal{L}_r$ the Lie algebra $\mathcal{L}$ with the new Lie bracket $[\cdot, \cdot]_r$.

Example 2.3. As an example, let us compute $[v^{ii}(m, n), v^{ii}(-n, -m)]_r$ for $1 \leq i \leq d$ and $m, n \in \mathbb{Z}_{>0}$ with $m \leq n$. By direct computation and (2.3), we see that in $\mathcal{L}$,

$$\begin{aligned} &[v^{ii}(m, n), v^{ii}(-n, -m)] \\ &= n(1 + \delta_{m,n})v^{ii}(-m, m) + m(1 + \delta_{m,n})v^{ii}(-n, n) + mn(1 + \delta_{m,n}). \end{aligned}$$

Hence, by the definition (2.4) of the Lie bracket $[\cdot, \cdot]_r$, we obtain

$$\begin{aligned}
& [v^{ii}(m, n), v^{ii}(-n, -m)]_r \\
&= \pi_1([v^{ii}(m, n), v^{ii}(-n, -m)]) + r\pi_2([v^{ii}(m, n), v^{ii}(-n, -m)]) \\
&= n(1 + \delta_{m,n})v^{ii}(-m, m) + m(1 + \delta_{m,n})v^{ii}(-n, n) + rmn(1 + \delta_{m,n}). \tag{2.5}
\end{aligned}$$

We now set

$$\begin{aligned}
\mathcal{B}_+ &:= \{v^{ij}(m, n) \in \mathcal{B} \mid m \in \mathbb{Z}_{\geq 0} \text{ or } n \in \mathbb{Z}_{\geq 0}\}, \\
\mathcal{B}_- &:= \{v^{ij}(m, n) \in \mathcal{B} \mid m, n \in \mathbb{Z}_{< 0}\},
\end{aligned}$$

and $\mathcal{L}_r^+ := (\text{Span}_{\mathbb{C}} \mathcal{B}_+) \oplus \mathbb{C}$. It is easily seen that $\mathcal{L}_r^+$ is a Lie subalgebra of $\mathcal{L}_r$. Let $\mathbb{C}\mathbf{1}$ be a one-dimensional $\mathcal{L}_r^+$ -module such that $x \cdot \mathbf{1} = 0$ for all $x \in \mathcal{B}_+$, and $s \cdot \mathbf{1} = s\mathbf{1}$ for each $s \in \mathbb{C} \subset \mathcal{L}_r^+$. Denote by M_r the $\mathcal{L}_r$ -module induced from the $\mathcal{L}_r^+$ -module $\mathbb{C}\mathbf{1}$, that is,

$$M_r := U(\mathcal{L}_r) \otimes_{U(\mathcal{L}_r^+)} \mathbb{C}\mathbf{1}.$$

Here we give a linear basis $\mathbb{B}$ of M_r by using the Poincaré-Birkhoff-Witt theorem. Take (and fix) a total ordering $\succ$ on the set $\mathcal{B}_-$. Denote by $\mathcal{S}$ the set of finite sequences of elements of $\mathcal{B}_-$ that is weakly decreasing with respect to the total ordering $\succ$. For $\mathbf{x} = (x_p \succeq x_{p-1} \succeq \cdots \succeq x_1) \in \mathcal{S}$ with $x_q \in \mathcal{B}_-$ for $1 \leq q \leq p$, we set

$$w(\mathbf{x}) := x_p x_{p-1} \cdots x_1 \mathbf{1} \in M_r.$$

In view of the Poincaré-Birkhoff-Witt theorem, $\mathbb{B} := \{w(\mathbf{x}) \mid \mathbf{x} \in \mathcal{S}\}$ is a linear basis of the $\mathcal{L}_r$ -module M_r .

Remark 2.4. We see from the definition (2.4) of the Lie bracket on $\mathcal{L}_r$ that $xy = yx$ for all $x, y \in \mathcal{B}_-$. Therefore, if $y_1, y_2, \dots, y_p \in \mathcal{B}_-$, then $y_1 y_2 \cdots y_p \mathbf{1} = w(\mathbf{x}) \in \mathbb{B}$, where $\mathbf{x} \in \mathcal{S}$ is the sequence of length p obtained by arranging $y_1, y_2, \dots, y_p$ in the weakly decreasing order with respect to the total ordering $\succ$. Also, we note that if $m, n \in \mathbb{Z}_{< 0}$, then $v^{ij}(m, n) = v^{ji}(n, m)$ for every $1 \leq i, j \leq d$ (see Remark 2.2).

If $\mathbf{x} = (x_p \succeq x_{p-1} \succeq \cdots \succeq x_1) \in \mathcal{S}$ with $x_q = v^{i_q j_q}(m_q, n_q) \in \mathcal{B}_-$ for $1 \leq q \leq p$, then we define the degree of $w(\mathbf{x}) \in \mathbb{B}$ by:

$$\deg(w(\mathbf{x})) = - \sum_{q=1}^p (m_q + n_q) \in \mathbb{Z}_{\geq 0}.$$

Then the $\mathcal{L}_r$ -module M_r admits the degree space decomposition as follows:

$$M_r = \bigoplus_{n \in \mathbb{Z}_{\geq 0}} (M_r)_n, \quad \text{where } (M_r)_n := \text{Span}_{\mathbb{C}} \{b \in \mathbb{B} \mid \deg b = n\}.$$

Note that

$$(M_r)_0 = \mathbb{C}\mathbf{1}, \quad \text{and} \quad (M_r)_1 = \{0\}. \quad (2.6)$$

Define an operator $L_r^{ij}(m) \in \text{End}(M_r)$ for $1 \leq i, j \leq d$ and $m \in \mathbb{Z}$ by:

$$L_r^{ij}(m) = \begin{cases} \frac{1}{2} \sum_{h \in \mathbb{Z}} v^{ij}(m-h, h) & \text{if } i \neq j \text{ or } m \neq 0, \\ \frac{1}{2} v^{ii}(0, 0) + \sum_{h \in \mathbb{Z}_{>0}} v^{ii}(-h, h) & \text{if } i = j \text{ and } m = 0, \end{cases} \quad (2.7)$$

and set

$$\omega_r^{ij} := L_r^{ij}(-2)\mathbf{1} \in (M_r)_2, \quad \text{and} \quad \omega := \sum_{i=1}^d \omega_r^{ii} \in (M_r)_2.$$

Remark that $L_r^{ij}(m) = L_r^{ji}(m)$ for every $1 \leq i, j \leq d$ and $m \in \mathbb{Z}$ (see Remark 2.2), and hence that $\omega_r^{ij} = \omega_r^{ji}$ for every $1 \leq i, j \leq d$. Let $\mathcal{J} := \{\omega_r^{ij}, \mathbf{1} \mid 1 \leq i, j \leq d\} \subset M_r$, and let $V_{\mathcal{J}}$ be the subspace of M_r spanned by all elements of the form: $L_r^{i_1 j_1}(m_1) L_r^{i_2 j_2}(m_2) \cdots L_r^{i_p j_p}(m_p) \mathbf{1}$ with $p \geq 0$, and $1 \leq i_q, j_q \leq d$, $m_q \in \mathbb{Z}$ for $1 \leq q \leq p$. Then, $V_{\mathcal{J}}$ also admits the degree space decomposition induced from that of M_r , i.e., $V_{\mathcal{J}} = \bigoplus_{n \in \mathbb{Z}_{\geq 0}} (V_{\mathcal{J}})_n$ with $(V_{\mathcal{J}})_n := V_{\mathcal{J}} \cap (M_r)_n$ for $n \in \mathbb{Z}_{\geq 0}$. We should remark that $(V_{\mathcal{J}})_0 = \mathbb{C}\mathbf{1}$ and $(V_{\mathcal{J}})_1 = \{0\}$ by (2.6).

Define a map $Y_0(\cdot, z) : \mathcal{J} \rightarrow \text{End}(V_{\mathcal{J}})[[z, z^{-1}]]$ by:

$$\begin{aligned} Y_0(\omega_r^{ij}, z) &= \sum_{m \in \mathbb{Z}} L_r^{ij}(m) z^{-m-2} \quad \text{for } 1 \leq i, j \leq d, \\ Y_0(\mathbf{1}, z) &= \text{id}_{V_{\mathcal{J}}}. \end{aligned} \quad (2.8)$$

The following theorem is the main result of [AM].

Theorem 2.5. *Keep the notation above. The map $Y_0(\cdot, z) : \mathcal{J} \rightarrow \text{End}(V_{\mathcal{J}})[[z, z^{-1}]]$ can be uniquely extended to a linear map $Y(\cdot, z) : V_{\mathcal{J}} \rightarrow \text{End}(V_{\mathcal{J}})[[z, z^{-1}]]$ in such a way that the quadruple $(V_{\mathcal{J}} = \bigoplus_{n \in \mathbb{Z}_{\geq 0}} (V_{\mathcal{J}})_n, Y(\cdot, z), \mathbf{1}, \omega)$ becomes a VOA of central charge dr , with $\mathbf{1}$ the vacuum element, and ω the Virasoro element. Furthermore, the Griess algebra of $V_{\mathcal{J}}$ is isomorphic to the Jordan algebra $\text{Sym}_d(\mathbb{C})$ of symmetric matrices.*

The purpose of this paper is to determine the condition of $r \in \mathbb{C}$ for the VOA $V_{\mathcal{J}}$ to be simple. The following theorem is the main result of this paper.

Theorem 2.6. *Keep the notation above. The VOA $V_{\mathcal{J}}$ is simple if and only if $r \in \mathbb{C}$ is not an integer, that is, $r \in \mathbb{C} \setminus \mathbb{Z}$.*

We will prove Theorem 2.6 as follows. First, in §3, we will show that $V_{\mathcal{J}} \subset M_r$ is, in fact, identical to the whole of M_r (Proposition 3.1), and then prove that the VOA $V_{\mathcal{J}}$ ($= M_r$) is simple if and only if M_r is irreducible as an $\mathcal{L}_r$ -module (Proposition 3.4). Let $\mathcal{L}_r^{(1)}$ be a Lie subalgebra of $\mathcal{L}_r$ generated by $\{v^{11}(m, n) \mid m, n \in \mathbb{Z} \text{ with } m \leq n\} \subset \mathcal{B}$, and set $M_r^{(1)} = U(\mathcal{L}_r^{(1)})\mathbf{1} \subset M_r$. In §4, it will be shown that the $\mathcal{L}_r$ -module M_r is irreducible if and only if $M_r^{(1)}$ is irreducible as an $\mathcal{L}_r^{(1)}$ -module (Proposition 4.1). In §5, we will prove that if $r \in \mathbb{C} \setminus \mathbb{Z}$, then $M_r^{(1)}$ is an irreducible $\mathcal{L}_r^{(1)}$ -module, and hence $V_{\mathcal{J}}$ is a simple VOA (Proposition 5.1). Finally, in §6, we will give some singular vectors of the $\mathcal{L}_r^{(1)}$ -module $M_r^{(1)}$ explicitly in the case that $r \in \mathbb{Z}$ (Proposition 6.1), which implies that $M_r^{(1)}$ is reducible, and hence $V_{\mathcal{J}}$ is not simple.

3 Simplicity of $V_{\mathcal{J}}$ and irreducibility of M_r .

3.1 Relation between $V_{\mathcal{J}}$ and M_r . As in the previous section, we fix $d \in \mathbb{Z}_{\geq 2}$ and $r \in \mathbb{C}$. This subsection is devoted to proving the following proposition.

Proposition 3.1. *The subspace $V_{\mathcal{J}} \subset M_r$ is identical to the whole of M_r , that is, $V_{\mathcal{J}} = M_r$ holds.*

In order to prove Proposition 3.1, we need some technical lemmas.

Lemma 3.2. (1) *For each $1 \leq i, j \leq d$, we have*

$$v^{ij}(-1, -1)\mathbf{1} = 2L_r^{ij}(-2)\mathbf{1}. \quad (3.1)$$

(2) *Let $1 \leq i, j \leq d$ with $i \neq j$, and $m, n \in \mathbb{Z}_{<0}$. Then,*

$$v^{ij}(m-1, n)\mathbf{1} = -\frac{1}{m}L_r^{ii}(-1)v^{ij}(m, n)\mathbf{1}. \quad (3.2)$$

(3) *Let $1 \leq i, j \leq d$ with $i \neq j$, and $m, n \in \mathbb{Z}_{<0}$. Then,*

$$v^{ii}(m-1, n)\mathbf{1} = \frac{2}{m(m-n+1)}L_r^{ii}(0)L_r^{ij}(-1)v^{ij}(n, m)\mathbf{1}. \quad (3.3)$$

Proof. (1) By the definition (2.7) of $L_r^{ij}(m)$,

$$2L_r^{ij}(-2)\mathbf{1} = \sum_{h \in \mathbb{Z}} v^{ij}(-2-h, h)\mathbf{1}.$$

Note that $v^{ij}(-2-h, h) = v^{ji}(h, -2-h)$ for all $h \in \mathbb{Z}$ (see Remark 2.2). Since $v^{ij}(m, n)\mathbf{1} = 0$ if $v^{ij}(m, n) \in \mathcal{B}_+$, it follows that $v^{ij}(-2-h, h)\mathbf{1} = 0$ unless $h = -1$. Thus we obtain $2L_r^{ij}(-2)\mathbf{1} = v^{ij}(-1, -1)\mathbf{1}$, and hence (3.1).

(2) As in the proof of part (1), we can easily show that

$$L_r^{ij}(-1)\mathbf{1} = \frac{1}{2} \sum_{h \in \mathbb{Z}} v^{ij}(-1-h, h)\mathbf{1} = 0 \quad (3.4)$$

for all $1 \leq i, j \leq d$. Hence,

$$\begin{aligned} L_r^{ii}(-1)v^{ij}(m, n)\mathbf{1} &= [L_r^{ii}(-1), v^{ij}(m, n)]\mathbf{1} \quad \text{since } L_r^{ii}(-1)\mathbf{1} = 0 \text{ by (3.4)} \\ &= \frac{1}{2} \sum_{h \in \mathbb{Z}} [v^{ii}(-1-h, h), v^{ij}(m, n)]\mathbf{1}. \end{aligned}$$

By direct computation (as in Example 2.3), we see that

$$\begin{aligned} [v^{ii}(-1-h, h), v^{ij}(m, n)]_r &= \\ \delta_{-1-h+m,0}(-1-h)v^{ij}(h, n) &+ \delta_{h+m,0}hv^{ij}(-1-h, n). \end{aligned}$$

Therefore,

$$\begin{aligned} &\frac{1}{2} \sum_{h \in \mathbb{Z}} [v^{ii}(-1-h, h), v^{ij}(m, n)]\mathbf{1} \\ &= \frac{1}{2} \sum_{h \in \mathbb{Z}} \{ \delta_{-1-h+m,0}(-1-h)v^{ij}(h, n) + \delta_{h+m,0}hv^{ij}(-1-h, n) \} \mathbf{1} \\ &= \frac{1}{2} \{ (-m)v^{ij}(m-1, n) + (-m)v^{ij}(m-1, n) \} \mathbf{1} \\ &= (-m)v^{ij}(m-1, n)\mathbf{1}. \end{aligned}$$

Thus we obtain $L_r^{ii}(-1)v^{ij}(m, n)\mathbf{1} = (-m)v^{ij}(m-1, n)$, and hence (3.2).

(3) We have

$$\begin{aligned} L_r^{ij}(-1)v^{ij}(n, m)\mathbf{1} &= [L_r^{ij}(-1), v^{ij}(n, m)]\mathbf{1} \quad \text{since } L_r^{ij}(-1)\mathbf{1} = 0 \text{ by (3.4)} \\ &= \frac{1}{2} \sum_{h \in \mathbb{Z}} [v^{ij}(-1-h, h), v^{ij}(n, m)]\mathbf{1}. \end{aligned}$$

As in Example 2.3, we see that

$$\begin{aligned} [v^{ij}(-1-h, h), v^{ij}(n, m)]_r &= \\ \delta_{h+m,0}hv^{ii}(n, -1-h) &+ \delta_{-1-h+n,0}(-1-h)v^{jj}(h, m). \end{aligned}$$

Therefore we get

$$\begin{aligned} &\frac{1}{2} \sum_{h \in \mathbb{Z}} [v^{ij}(-1-h, h), v^{ij}(n, m)]\mathbf{1} \\ &= \frac{1}{2} \sum_{h \in \mathbb{Z}} \{ \delta_{h+m,0}hv^{ii}(n, -1-h) + \delta_{-1-h+n,0}(-1-h)v^{jj}(h, m) \} \mathbf{1} \\ &= \frac{1}{2} \{ (-m)v^{ii}(n, m-1)\mathbf{1} + (-n)v^{jj}(n-1, m)\mathbf{1} \}, \end{aligned}$$

and hence

$$L_r^{ij}(-1)v^{ij}(n, m)\mathbf{1} = -\frac{m}{2}v^{ii}(n, m-1)\mathbf{1} - \frac{n}{2}v^{jj}(n-1, m)\mathbf{1}.$$

Now, since $L_r^{ii}(0)\mathbf{1} = 0$, it follows that

$$\begin{aligned} L_r^{ii}(0)L_r^{ij}(-1)v^{ij}(n, m)\mathbf{1} \\ &= -\frac{m}{2}L_r^{ii}(0)v^{ii}(n, m-1)\mathbf{1} - \frac{n}{2}L_r^{ii}(0)v^{jj}(n-1, m)\mathbf{1} \\ &= -\frac{m}{2}[L_r^{ii}(0), v^{ii}(n, m-1)]\mathbf{1} - \frac{n}{2}[L_r^{ii}(0), v^{jj}(n-1, m)]\mathbf{1}. \end{aligned}$$

It can be easily checked that $[L_r^{ii}(0), v^{jj}(n-1, m)] = 0$ since $i \neq j$. Thus,

$$\begin{aligned} &-\frac{m}{2}[L_r^{ii}(0), v^{ii}(n, m-1)]\mathbf{1} - \frac{n}{2}[L_r^{ii}(0), v^{jj}(n-1, m)]\mathbf{1} \\ &= -\frac{m}{2}[L_r^{ii}(0), v^{ii}(n, m-1)]\mathbf{1} \\ &= \left\{ -\frac{m}{4} \underbrace{[v^{ii}(0, 0), v^{ii}(n, m-1)]}_{=0} - \frac{m}{2} \sum_{h \in \mathbb{Z}_{>0}} [v^{ii}(-h, h), v^{ii}(n, m-1)] \right\} \mathbf{1}. \end{aligned}$$

Since $n, m-1 \in \mathbb{Z}_{<0}$, it follows that

$$[v^{ii}(-h, h), v^{ii}(n, m-1)]_r = \delta_{h+n,0} h v^{ii}(-h, m-1) + \delta_{h+m-1,0} h v^{ii}(n, -h)$$

for $h \in \mathbb{Z}_{\geq 1}$. Therefore,

$$\begin{aligned} &-\frac{m}{2} \sum_{h \in \mathbb{Z}_{>0}} [v^{ii}(-h, h), v^{ii}(n, m-1)]\mathbf{1} \\ &= -\frac{m}{2} \sum_{h \in \mathbb{Z}_{>0}} \{ \delta_{h+n,0} h v^{ii}(-h, m-1) + \delta_{h+m-1,0} h v^{ii}(n, -h) \} \mathbf{1} \\ &= -\frac{m}{2} \{ (-n)v^{ii}(n, m-1) + (-m+1)v^{ii}(n, m-1) \} \mathbf{1} \\ &= \frac{m(m+n-1)}{2} v^{ii}(m-1, n)\mathbf{1}. \end{aligned}$$

Thus we obtain

$$L_r^{ii}(0)L_r^{ij}(-1)v^{ij}(n, m)\mathbf{1} = \frac{m(m+n-1)}{2} v^{ii}(m-1, n)\mathbf{1},$$

and hence (3.3). This completes the proof of the lemma. $\square$

Lemma 3.3. (1) Let $1 \leq i, j \leq d$ with $i \neq j$, and $m, n \in \mathbb{Z}_{<0}$. Then,

$$v^{ij}(m, n)\mathbf{1} = \alpha L_r^{ii}(-1)^{-m-1} L_r^{jj}(-1)^{-n-1} L_r^{ij}(-2)\mathbf{1} \quad (3.5)$$

for some $\alpha \in \mathbb{C} \setminus \{0\}$.

(2) Let $1 \leq i \leq d$, and $m, n \in \mathbb{Z}_{<0}$ with $m \leq -2$. Take $1 \leq j \leq d$ with $j \neq i$ arbitrarily. Then,

$$v^{ii}(m, n)\mathbf{1} = \beta L_r^{ii}(0)L_r^{ij}(-1)L_r^{ii}(-1)^{-n-1}L_r^{jj}(-1)^{-m-2}L_r^{ij}(-2)\mathbf{1} \quad (3.6)$$

for some $\beta \in \mathbb{C} \setminus \{0\}$.

Proof. (1) We prove (3.5) by induction on $-m-n$ (note that $-m-n \geq 2$). If $-m-n = 2$, that is, $m = n = -1$, then (3.5) follows immediately from (3.1). Assume that $-m-n > 2$. By Remark 2.2, we may assume that $m < -1$. Then, by (3.2), we have

$$v^{ij}(m, n)\mathbf{1} = -\frac{1}{m+1}L_r^{ii}(-1)v^{ij}(m+1, n)\mathbf{1}.$$

Applying the inductive assumption to the right-hand side of the equation above, we obtain

$$\begin{aligned} v^{ij}(m, n)\mathbf{1} &= -\frac{1}{m+1}L_r^{ii}(-1)v^{ij}(m+1, n)\mathbf{1} \\ &= -\frac{1}{m+1}L_r^{ii}(-1)\{\alpha L_r^{ii}(-1)^{-m-2}L_r^{jj}(-1)^{-n-1}L_r^{ij}(-2)\mathbf{1}\} \\ &= -\frac{\alpha}{m+1}L_r^{ii}(-1)^{-m-1}L_r^{jj}(-1)^{-n-1}L_r^{ij}(-2)\mathbf{1}, \end{aligned}$$

where $\alpha \in \mathbb{C} \setminus \{0\}$. Thus we have proved part (1).

(2) Using (3.3) and (3.5), we have

$$\begin{aligned} v^{ii}(m, n)\mathbf{1} &= \frac{2}{(m+1)(m+n)}L_r^{ii}(0)L_r^{ij}(-1)v^{ij}(n, m+1)\mathbf{1} \quad \text{by (3.3)} \\ &= \frac{2\beta}{(m+1)(m+n)}L_r^{ii}(0)L_r^{ij}(-1)L_r^{ii}(-1)^{-n-1}L_r^{jj}(-1)^{-m-2}L_r^{ij}(-2)\mathbf{1} \quad \text{by (3.6),} \end{aligned}$$

where $\beta \in \mathbb{C} \setminus \{0\}$. Thus we have proved part (2), thereby completing the proof of Lemma 3.3. $\square$

Proof of Proposition 3.1. We set

$$U := \text{Span}_{\mathbb{C}} \left\{ L_r^{i_1 j_1}(m_1)L_r^{i_2 j_2}(m_2) \cdots L_r^{i_p j_p}(m_p)\mathbf{1} \mid \begin{array}{l} p \geq 0, \text{ and } 1 \leq i_q, j_q \leq d, \\ m_q \in \{-2, -1, 0\} \text{ for } 1 \leq q \leq p \end{array} \right\}.$$

In order to prove Proposition 3.1, it suffices to show that $U = M_r$. Indeed, it is obvious from the definition of $V_{\mathcal{J}}$ that $U \subset V_{\mathcal{J}}$. Hence, if $U = M_r$ holds, then $M_r = U \subset V_{\mathcal{J}} \subset M_r$, which implies that $V_{\mathcal{J}} = M_r$.

Claim. We have $xU \subset U$ for every $x \in \mathcal{B}$.

Proof of Claim. Fix $x \in \mathcal{B}$. It suffices to prove that

$$xL_r^{i_1j_1}(m_1)L_r^{i_2j_2}(m_2) \cdots L_r^{i_pj_p}(m_p)\mathbf{1} \in U$$

for every $p \geq 0$ and every $1 \leq i_q, j_q \leq d$, $m_q \in \{-2, -1, 0\}$ for $1 \leq q \leq p$. We prove this by induction on p . It is obvious that $x\mathbf{1} = 0 \in U$ if $x \in \mathcal{B}_+$. Also, we see from equations (3.5) and (3.6) that $x\mathbf{1}$ is contained in U if $x \in \mathcal{B}_-$. Thus the assertion holds when $p = 0$. Assume that $p > 0$. Then we have

$$\begin{aligned} & xL_r^{i_1j_1}(m_1)L_r^{i_2j_2}(m_2) \cdots L_r^{i_pj_p}(m_p)\mathbf{1} = \\ & [x, L_r^{i_1j_1}(m_1)]L_r^{i_2j_2}(m_2) \cdots L_r^{i_pj_p}(m_p)\mathbf{1} + L_r^{i_1j_1}(m_1)\{xL_r^{i_2j_2}(m_2) \cdots L_r^{i_pj_p}(m_p)\mathbf{1}\}. \end{aligned} \quad (3.7)$$

Since $xL_r^{i_2j_2}(m_2) \cdots L_r^{i_pj_p}(m_p)\mathbf{1}$ is contained in U by the inductive assumption, and since $m_1 \in \{-2, -1, 0\}$, it follows that the second term of the right-hand side of (3.7) is contained in U . Now we deduce from the definition of the Lie bracket on $\mathcal{L}_r$ and the definition (2.7) of $L_r^{ij}(m)$ that $[x, L_r^{i_1j_1}(m_1)]$ can be written in the form:

$$[x, L_r^{i_1j_1}(m_1)] = \alpha_1 y_1 + \alpha_2 y_2 + \cdots + \alpha_s y_s + \beta$$

for some $\alpha_1, \alpha_2, \dots, \alpha_s, \beta \in \mathbb{C}$ and $y_1, y_2, \dots, y_s \in \mathcal{B}$. By substituting this into the first term of the right-hand side of (3.7), we see that

$$\begin{aligned} & [x, L_r^{i_1j_1}(m_1)]L_r^{i_2j_2}(m_2) \cdots L_r^{i_pj_p}(m_p)\mathbf{1} \\ &= \sum_{1 \leq s' \leq s} \alpha_{s'} \underbrace{y_{s'} L_r^{i_2j_2}(m_2) \cdots L_r^{i_pj_p}(m_p)\mathbf{1}}_{\in U \text{ by the inductive assumption}} + \beta \underbrace{L_r^{i_2j_2}(m_2) \cdots L_r^{i_pj_p}(m_p)\mathbf{1}}_{\in U}, \end{aligned}$$

and hence that the first term also is contained in U . Therefore we conclude that the left-hand side of (3.7) is contained in U , thereby completing the proof of Claim.

The claim above implies that U is an $\mathcal{L}_r$ -submodule of M_r which contains $\mathbf{1}$. Hence we conclude that $U = M_r$, thereby completing the proof of Proposition 3.1. $\square$

3.2 Relation between the simplicity of $V_{\mathcal{J}}$ and the irreducibility of M_r . In this subsection, we prove the following proposition.

Proposition 3.4. *The VOA $V_{\mathcal{J}}$ ($= M_r$) is simple if and only if M_r is irreducible as an $\mathcal{L}_r$ -module.*

First let us show the following lemma, needed in the proof of Proposition 3.4.

Lemma 3.5. *Let $1 \leq i, j \leq d$ with $i \neq j$, and $m, n \in \mathbb{Z}_{<0}$. The vertex operator $Y(v^{ij}(m, n)\mathbf{1}, z)$ of $v^{ij}(m, n)\mathbf{1} \in M_r = V_{\mathcal{J}}$ is given by:*

$$Y(v^{ij}(m, n)\mathbf{1}, z) = (-1)^{-m-n} \times \sum_{l \in \mathbb{Z}} \left\{ \sum_{k \in \mathbb{Z}} \binom{l+n-k}{-m-1} \binom{k-n-1}{-n-1} v^{ij}(l+m+n+1-k, k) \right\} z^{-l-1}. \quad (3.8)$$

Proof. We prove the lemma by induction on $-m-n$ (note that $-m-n \geq 2$). If $-m-n = 2$, that is, $m = n = -1$, then it follows that

$$\begin{aligned} Y(v^{ij}(-1, -1)\mathbf{1}, z) &= 2Y(L_r^{ij}(-2)\mathbf{1}, z) \quad \text{by (3.1)} \\ &= 2Y(\omega_r^{ij}, z) = 2 \sum_{l \in \mathbb{Z}} L_r^{ij}(l) z^{-l-2} \quad \text{by (2.8) and Theorem 2.5} \\ &= \sum_{l \in \mathbb{Z}} \left\{ \sum_{k \in \mathbb{Z}} v^{ij}(l-k, k) \right\} z^{-l-2} = \sum_{l \in \mathbb{Z}} \left\{ \sum_{k \in \mathbb{Z}} v^{ij}(l-1-k, k) \right\} z^{-l-1}. \end{aligned}$$

Therefore the equation (3.8) holds if $-m-n = 2$. Assume that $-m-n > 2$. By Remark 2.2, we may assume that $m < -1$. It follows from (3.2) that

$$Y(v^{ij}(m, n)\mathbf{1}, z) = \frac{1}{-m-1} Y(L_r^{ii}(-1)v^{ij}(m+1, n)\mathbf{1}, z).$$

By using the commutator formula (see [LL, p. 54]), we deduce that

$$\begin{aligned} (L_r^{ii}(-1)v^{ij}(m+1, n)\mathbf{1})_l &= ((\omega_r^{ii})_0 v^{ij}(m+1, n)\mathbf{1})_l = [(\omega_r^{ii})_0, (v^{ij}(m+1, n)\mathbf{1})_l] \\ &= [L_r^{ii}(-1), (v^{ij}(m+1, n)\mathbf{1})_l]. \end{aligned}$$

Also, it follows from the inductive assumption that

$$(v^{ij}(m+1, n)\mathbf{1})_l = (-1)^{-m-1-n} \times \left\{ \sum_{k \in \mathbb{Z}} \binom{l+n-k}{-m-2} \binom{k-n-1}{-n-1} v^{ij}(l+m+n+2-k, k) \right\}.$$

Combining these equations, we get

$$\begin{aligned} Y(v^{ij}(m, n)\mathbf{1}, z) &= \frac{(-1)^{-m-1-n}}{-m-1} \sum_{l \in \mathbb{Z}} \left\{ \sum_{k \in \mathbb{Z}} \binom{l+n-k}{-m-2} \binom{k-n-1}{-n-1} \times \right. \\ &\quad \left. [L_r^{ii}(-1), v^{ij}(l+m+n+2-k, k)] \right\} z^{-l-1}. \end{aligned}$$

It can be easily checked by direct computation that

$$\begin{aligned}
& [L_r^{ii}(-1), v^{ij}(l+m+n+2-k, k)] \\
&= \frac{1}{2} \sum_{h \in \mathbb{Z}} [v^{ii}(-1-h, h), v^{ij}(l+m+n+2-k, k)] \\
&= -(l+m+n+2-k) v^{ij}(l+m+n+1-k, k).
\end{aligned}$$

Thus we obtain

$$\begin{aligned}
& Y(v^{ij}(m, n)\mathbf{1}, z) \\
&= (-1)^{-m-n} \sum_{l \in \mathbb{Z}} \left\{ \sum_{k \in \mathbb{Z}} \frac{1}{-m-1} \binom{l+n-k}{-m-2} (l+m+n+2-k) \times \right. \\
&\quad \left. \binom{k-n-1}{-n-1} v^{ij}(l+m+n+1-k, k) \right\} z^{-l-1} \\
&= (-1)^{-m-n} \sum_{l \in \mathbb{Z}} \left\{ \sum_{k \in \mathbb{Z}} \binom{l+n-k}{-m-1} \binom{k-n-1}{-n-1} v^{ij}(l+m+n+1-k, k) \right\} z^{-l-1}.
\end{aligned}$$

Thus we have proved Lemma 3.5. $\square$

Proof of Proposition 3.4. First, we show the “if” part. Assume that $V_{\mathcal{J}} = M_r$ is not simple, and let $W \subset V_{\mathcal{J}} = M_r$ be a proper ideal of the VOA $V_{\mathcal{J}}$. Let us show that W is a (proper) $\mathcal{L}_r$ -submodule of M_r .

Claim 1. *Let $1 \leq i, j \leq d$ with $i \neq j$. Then, $v^{ij}(m, n)W \subset W$ for all $m, n \in \mathbb{Z}$.*

Proof of Claim 1. Let $u \in W$. We prove that $v^{ij}(m, n)u \in W$ for all $m, n \in \mathbb{Z}$, which is equivalent to showing that $v^{ij}(s-t, t)u \in W$ for all $s, t \in \mathbb{Z}$. Fix $s \in \mathbb{Z}$. If $v^{ij}(s-t, t)u = 0$ for all $t \in \mathbb{Z}$, then the assertion is obvious. So, let us assume that $v^{ij}(s-t, t)u \neq 0$ for some $t \in \mathbb{Z}$. Take $t_1, t_2 \in \mathbb{Z}$ with $t_1 \leq t_2$ in such a way that if $v^{ij}(s-t, t)u \neq 0$, then $t_1 \leq t \leq t_2$. By Lemma 3.5, we see that

$$\begin{aligned}
(v^{ij}(-p, -1)\mathbf{1})_{s+p} u &= (-1)^{p+1} \sum_{t \in \mathbb{Z}} \binom{s+p-1-t}{p-1} v^{ij}(s-t, t)u \\
&= \sum_{t_1 \leq t \leq t_2} (-1)^{p+1} \binom{s+p-1-t}{p-1} v^{ij}(s-t, t)u.
\end{aligned}$$

for $p > 0$. Because W is an ideal of the VOA $V_{\mathcal{J}} = M_r$, it follows that

$$(v^{ij}(-p, -1)\mathbf{1})_{s+p} u \in W$$

for all $p > 0$. Hence, in order to prove that $v^{ij}(s-t, t)u \in W$ for all $t_1 \leq t \leq t_2$, it suffices to show that the matrix

$$\left((-1)^{p+1} \binom{s+p-1-t}{p-1} \right)_{\substack{1 \leq p \leq t_2-t_1+1 \\ t_1 \leq t \leq t_2}} \quad (3.9)$$

is invertible. So we show the following: Let $L \in \mathbb{Z}$, and $M \in \mathbb{Z}_{>0}$. Then,

$$\det(a_{p,N})_{1 \leq p, N \leq M} \neq 0, \quad \text{where } a_{p,N} := \binom{L+p-N}{p-1}. \quad (3.10)$$

We prove (3.10) by induction on the size M of the matrix. The claim is obvious when $M = 1$. Assume that $M > 1$. Using the formula

$$a_{p,N} - a_{p,N-1} = -\binom{L+p-N}{p-2} \quad \text{for } 2 \leq N \leq M,$$

we deduce that

$$\begin{aligned} \det(a_{p,N})_{1 \leq p, N \leq M} &= \det \left(-\binom{L+p-N}{p-2} \right)_{2 \leq p, N \leq M} \\ &= (-1)^{M-1} \det \left(\binom{L+p-N}{p-2} \right)_{2 \leq p, N \leq M} \\ &= (-1)^{M-1} \underbrace{\det(a_{p',N'})_{1 \leq p', N' \leq M-1}}_{\neq 0 \text{ by the inductive assumption}}, \end{aligned}$$

where $p' := p-1$ and $N' := N-1$. Thus we obtain (3.10). We see from (3.10) with $L = s-t_1$, $M = t_2-t_1+1$, and $N = t-t_1+1$ that the determinant of the matrix (3.9) is not equal to 0, and hence the matrix (3.9) is invertible. Thus we have proved Claim 1.

Claim 2. *Let $1 \leq i \leq d$. Then, $v^{ii}(m, n)W \subset W$ for all $m, n \in \mathbb{Z}$.*

Proof of Claim 2. Let $u \in W$. Take $1 \leq j \leq d$ with $j \neq i$ arbitrarily, and take $N \in \mathbb{Z}_{<0}$ in such a way that $v^{jj}(N, -N)u = 0$. By direct computation, we see that

$$\begin{aligned} [v^{ij}(m, -N), v^{ij}(n, N)]u &= -Nv^{ii}(m, n)u + \delta_{m+n,0}mv^{jj}(N, -N)u + \alpha u \\ &= -Nv^{ii}(m, n)u + \alpha u \quad \text{since } v^{jj}(N, -N)u = 0 \end{aligned}$$

for some $\alpha \in \mathbb{C}$. Since $[v^{ij}(m, -N), v^{ij}(n, N)]u$ is contained in W by Claim 1, we conclude that $v^{ii}(m, n)u \in W$, thereby completing the proof of Claim 2.

It follows from Claims 1 and 2 that $v^{ij}(m, n)W \subset W$ for all $1 \leq i, j \leq d$ and $m, n \in \mathbb{Z}$, which implies that W is an $\mathcal{L}_r$ -submodule of M_r . Thus we have proved the “if” part of Proposition 3.4.

Next, we show the “only if” part. Assume that $M_r = V_{\mathcal{J}}$ is a reducible $\mathcal{L}_r$ -module, and let $W \subset M_r = V_{\mathcal{J}}$ be a proper $\mathcal{L}_r$ -submodule. Let $v \in V_{\mathcal{J}}$, and $l \in \mathbb{Z}$. By the definition of the vertex operator of $V_{\mathcal{J}}$, we deduce that $v_l \in \text{End}(V_{\mathcal{J}})$ can be written as an (infinite) linear combination of products of $v^{ij}(m, n)$, $1 \leq i, j \leq d$, $m, n \in \mathbb{Z}$. Since $v^{ij}(m, n)W \subset W$ by assumption, we have $v_l W \subset W$. Therefore, W is a proper ideal of the VOA $V_{\mathcal{J}}$, and hence $V_{\mathcal{J}}$ is not simple. This completes the proof of Proposition 3.4. $\square$

The next corollary follows immediately from the proof of Proposition 3.4.

Corollary 3.6. *Assume that the VOA $V_{\mathcal{J}}$ is not simple, or equivalently, the $\mathcal{L}_r$ -module M_r is reducible. Then, $W \subset V_{\mathcal{J}} (= M_r)$ is the maximal proper ideal of the VOA $V_{\mathcal{J}}$ if and only if W is the maximal proper $\mathcal{L}_r$ -submodule of M_r .*

4 Irreducibility of M_r and $M_r^{(1)}$.

Let $\mathcal{L}_r^{(1)}$ be the Lie subalgebra of $\mathcal{L}_r$ generated by $\mathcal{B}^{(1)} := \{v^{11}(m, n) \mid m, n \in \mathbb{Z} \text{ with } m \leq n\} \subset \mathcal{B}$. Set $M_r^{(1)} = U(\mathcal{L}_r^{(1)})\mathbf{1} \subset M_r$. In this section, we prove the following proposition.

Proposition 4.1. *The $\mathcal{L}_r$ -module M_r is irreducible if and only if $M_r^{(1)}$ is irreducible as an $\mathcal{L}_r^{(1)}$ -module.*

4.1 Proof of the “if” part of Proposition 4.1. In this subsection, we show that if $M_r^{(1)}$ is an irreducible $\mathcal{L}_r^{(1)}$ -module, then M_r is an irreducible $\mathcal{L}_r$ -module. This assertion follows immediately from the next lemma.

Lemma 4.2. *Let W be a nonzero $\mathcal{L}_r$ -submodule of M_r . Then, $W \cap M_r^{(1)}$ is a nonzero $\mathcal{L}_r^{(1)}$ -submodule of $M_r^{(1)}$.*

Indeed, if $W \subset M_r$ is a nonzero $\mathcal{L}_r$ -submodule of M_r , then it follows from Lemma 4.2 that $W \cap M_r^{(1)}$ is a nonzero $\mathcal{L}_r^{(1)}$ -submodule of $M_r^{(1)}$. Since $M_r^{(1)}$ is assumed to be an irreducible $\mathcal{L}_r^{(1)}$ -module, we have $W \cap M_r^{(1)} = M_r^{(1)}$. In particular, W contains $\mathbf{1} \in M_r$, which implies that $W = M_r$.

In order to prove Lemma 4.2, we introduce a weight space decomposition of M_r . Define

$$\mathcal{H} := \bigoplus_{1 \leq k \leq d, l \in \mathbb{Z}_{<0}} \mathbb{C} v^{kk}(l, -l) \subset \mathcal{L}_r.$$

Then it can be easily seen that $\mathcal{H}$ is an abelian Lie subalgebra of $\mathcal{L}_r$. Set $h_{k,l} := -(1/l) v^{kk}(l, -l) \in \mathcal{H}$ for $1 \leq k \leq d$ and $l \in \mathbb{Z}_{<0}$. By simple computation, we see that for $1 \leq k \leq d$ and $l \in \mathbb{Z}_{<0}$, $1 \leq i, j \leq d$ and $m, n \in \mathbb{Z}$,

$$[h_{k,l}, v^{ij}(m, n)]_r = (\delta_{(k,l), (i,m)} + \delta_{(k,l), (j,n)} - \delta_{(k,-l), (i,m)} - \delta_{(k,-l), (j,n)}) v^{ij}(m, n). \quad (4.1)$$

Let $\Lambda_{k,l} \in \mathcal{H}^* := \text{Hom}_{\mathbb{C}}(\mathcal{H}, \mathbb{C})$ be the dual basis of $h_{k,l} \in \mathcal{H}$ for $1 \leq k \leq d$ and $l \in \mathbb{Z}_{<0}$, and set

$$Q_+ := \sum_{1 \leq k \leq d, l \in \mathbb{Z}_{<0}} \mathbb{Z}_{\geq 0} \Lambda_{k,l} \subset \mathcal{H}^*.$$

We see from (4.1) that for each $\mathbf{x} \in \mathcal{S}$, the basis element $w(\mathbf{x})$ of M_r is contained in the “weight space” $(M_r)^\lambda := \{u \in M_r \mid hu = \lambda(h)u \text{ for all } h \in \mathcal{H}\}$ of weight λ for some $\lambda \in Q_+$ (see also Remark 4.4 below). Thus the $\mathcal{L}_r$ -module M_r admits the weight space decomposition with respect to the abelian Lie subalgebra $\mathcal{H} \subset \mathcal{L}_r$ as:

$$M_r = \bigoplus_{\lambda \in Q_+} (M_r)^\lambda. \quad (4.2)$$

Remark 4.3. Let $1 \leq i, j \leq d$, and let $m, n \in \mathbb{Z}$. It follows from (4.1) that $v^{ij}(m, n) \in \text{End}_{\mathbb{C}}(M_r)$ is a homogeneous operator of weight $-\text{sign}(m)(\Lambda_{i,-m} + \Lambda_{i,m}) - \text{sign}(n)(\Lambda_{j,-n} + \Lambda_{j,n})$, where for $N \in \mathbb{Z}$,

$$\text{sign}(N) := \begin{cases} 1 & \text{if } N > 0, \\ 0 & \text{if } N = 0, \\ -1 & \text{if } N < 0, \end{cases}$$

and for convenience, $\Lambda_{k,l} := 0$ for all $1 \leq k \leq d$ and $l \in \mathbb{Z}_{\geq 0}$.

Remark 4.4. Let $\mathbf{x} = (x_p \succeq x_{p-1} \succeq \cdots \succeq x_1) \in \mathcal{S}$ with $x_q = v^{i_q j_q}(m_q, n_q) \in \mathcal{B}_-$ for $1 \leq q \leq p$. For $1 \leq k \leq d$ and $l \in \mathbb{Z}_{<0}$, we define

$$\begin{aligned} \nu(\mathbf{x}, (k, l)) &:= \\ &\# \{1 \leq q \leq p \mid (i_q, m_q) = (k, l)\} + \# \{1 \leq q \leq p \mid (j_q, n_q) = (k, l)\}. \end{aligned}$$

Namely, $\nu(\mathbf{x}, (k, l))$ denotes “the number of $v^k(l)$ appearing in $\mathbf{x}$ ” (see (2.2)). Then we deduce from Remark 4.3 that the weight of $w(\mathbf{x})$ is equal to

$$\sum_{1 \leq k \leq d, l \in \mathbb{Z}_{<0}} \nu(\mathbf{x}, (k, l)) \Lambda_{k,l} \in Q_+.$$

Proof of Lemma 4.2. Remark that the submodule $W \subset M_r$ also admits the weight space decomposition $W = \bigoplus_{\lambda \in Q_+} W^\lambda$, where $W^\lambda := W \cap (M_r)^\lambda$. Let u be a nonzero homogeneous element of W , that is, $u \in W^\lambda \setminus \{0\}$ for some $\lambda \in Q_+$. It suffices to show that there exists $x \in U(\mathcal{L}_r)$ such that $xu \in M_r^{(1)} \setminus \{0\}$. For each $\xi \in Q_+$, we set

$$\theta(\xi) := \sum_{2 \leq k \leq d, l \in \mathbb{Z}_{<0}} \xi(h_{k,l}). \quad (4.3)$$

We show the claim above by induction on $\theta(\lambda)$. If $\theta(\lambda) = 0$, then the claim is obvious since $u \in M_r^{(1)}$ (see Remark 4.4). Assume that $\theta(\lambda) > 0$. Let $2 \leq i \leq d$ and $m \in \mathbb{Z}_{<0}$

be such that $\lambda(h_{i,m}) \geq 1$, and let $N \in \mathbb{Z}_{<0}$ be such that $\lambda(h_{1,N}) = 0$. Then we have $v^{1i}(-N, m)v^{1i}(N, -m)u \neq 0$. Indeed, since

$$[v^{1i}(-N, m), v^{1i}(N, -m)]_r = (-N)v^{ii}(m, -m) + mv^{11}(N, -N)$$

by direct computation, it follows that

$$\begin{aligned} v^{1i}(-N, m)v^{1i}(N, -m)u &= \{(-N)v^{ii}(m, -m) + mv^{11}(N, -N)\}u \\ &\quad + v^{1i}(N, -m)v^{1i}(-N, m)u. \end{aligned}$$

Here we note that $\lambda + \Lambda_{i,m} - \Lambda_{1,N}$ is not contained in Q_+ since $\lambda(h_{1,N}) = 0$. Because the weight of $v^{1i}(-N, m)u$ is equal to $\lambda + \Lambda_{i,m} - \Lambda_{1,N} \notin Q_+$ by Remark 4.3, we see from (4.2) that $v^{1i}(-N, m)u = 0$. Hence we get

$$\begin{aligned} v^{1i}(-N, m)v^{1i}(N, -m)u &= \{(-N)v^{ii}(m, -m) + mv^{11}(N, -N)\}u \\ &= mN \underbrace{\lambda(h_{i,m})}_{\geq 1} u - mN \underbrace{\lambda(h_{1,N})}_{=0} u \neq 0. \end{aligned}$$

Thus we obtain $v^{1i}(-N, m)v^{1i}(N, -m)u \neq 0$, which implies that $v^{1i}(N, -m)u \neq 0$. It follows from Remark 4.3 that the weight λ' of $v^{1i}(N, -m)u$ is equal to $\lambda - \Lambda_{i,m} + \Lambda_{1,N}$. Since $\theta(\lambda') = \theta(\lambda) - 1$, there exists $x' \in U(\mathcal{L}_r)$ such that $x'v^{1i}(N, -m)u \in M_r^{(1)}$ by the inductive assumption. Thus we have proved Lemma 4.2. $\square$

4.2 Proof of the “only if” part of Proposition 4.1. Assume that $W \subset M_r^{(1)}$ is an $\mathcal{L}_r^{(1)}$ -submodule of $M_r^{(1)}$. Then we see that W is stable under the action of $\mathcal{H}$; indeed, we have $h_{1,l}W \subset W$ for all $l \in \mathbb{Z}_{<0}$ by assumption, and $h_{k,l}W = \{0\}$ for all $2 \leq k \leq d$ and $l \in \mathbb{Z}_{<0}$ (see (4.1) and Remark 4.4). Thus, W also admits the weight space decomposition as follows: $W = \bigoplus_{\lambda \in Q_+} W^\lambda$ with $W^\lambda = W \cap (M_r)^\lambda$. Since $W \subset M_r^{(1)} = U(\mathcal{L}_r^{(1)})\mathbf{1}$, we deduce from Remark 4.3 (see also Remark 4.4) that $W^\lambda = \{0\}$ unless $\lambda \in Q_+^{(1)} := \sum_{l \in \mathbb{Z}_{<0}} \mathbb{Z}_{\geq 0} \Lambda_{1,l}$. Hence we have

$$W = \bigoplus_{\lambda \in Q_+} W^\lambda = \bigoplus_{\lambda \in Q_+^{(1)}} W^\lambda. \quad (4.4)$$

Now, let us prove that if M_r is an irreducible $\mathcal{L}_r$ -module, then $M_r^{(1)}$ is an irreducible $\mathcal{L}_r^{(1)}$ -module. It suffices to show the next lemma.

Lemma 4.5. *Assume that $M_r^{(1)}$ is a reducible $\mathcal{L}_r^{(1)}$ -module, and let $W \subset M_r^{(1)}$ be a proper $\mathcal{L}_r^{(1)}$ -submodule of $M_r^{(1)}$. Let u be a nonzero homogeneous element of W , that is, $u \in W^\lambda \setminus \{0\}$ for some $\lambda \in Q_+^{(1)}$. Then the $\mathcal{L}_r$ -module $U(\mathcal{L}_r)u \subset M_r$ generated by the u is a proper submodule of M_r . Therefore the $\mathcal{L}_r$ -module M_r is reducible.*

Proof. Suppose that $U(\mathcal{L}_r)u$ coincides with the whole of M_r . Then there exists $x \in U(\mathcal{L}_r)$ such that $xu \in \mathbb{C}\mathbf{1} \setminus \{0\}$. Let

$$\mathcal{B}_1 := \mathcal{B}_- \cup \{v^{1j}(m, n) \in \mathcal{B} \mid 2 \leq j \leq d, m \in \mathbb{Z}, n \in \mathbb{Z}_{<0}\}, \quad \mathcal{B}_2 := \mathcal{B} \setminus \mathcal{B}_1.$$

Since an element in $\mathbb{C} \subset \mathcal{L}_r$ acts as a scalar multiple, we may assume, by the Poincaré-Birkhoff-Witt theorem, that the $x \in U(\mathcal{L}_r)$ above is of the form: $x = \sum_{1 \leq t \leq s} \alpha_t y_t z_t$, where y_t (resp., z_t) is a product of elements in $\mathcal{B}_1$ (resp., $\mathcal{B}_2$) for each $1 \leq t \leq s$, and $\alpha_t \in \mathbb{C}$ for each $1 \leq t \leq s$. Because u is a homogeneous element, we see from Remark 4.3 that $y_t z_t u$ are also homogeneous elements for all $1 \leq t \leq s$. Since $xu \in \mathbb{C}\mathbf{1} \setminus \{0\}$, and since $(M_r)^0 = (M_r)_0 = \mathbb{C}\mathbf{1}$, it follows that $y_t z_t u \in \mathbb{C}\mathbf{1} \setminus \{0\}$ for some $1 \leq t \leq s$. Thus we may assume from the beginning that x is of the form: $x = yz$, where y (resp., z) is a product of elements in $\mathcal{B}_1$ (resp., $\mathcal{B}_2$).

Suppose that $y \neq 1$. Because $xu = yzu \in \mathbb{C}\mathbf{1} \setminus \{0\} \subset (M_r)^0$, we deduce from the definition of the set $\mathcal{B}_1$ and Remark 4.3 that the weight of zu is not contained in Q_+ . Hence, $zu = 0$ by (4.2), which is a contradiction. Thus we get $y = 1$. Write $z = z_p z_{p-1} \cdots z_1$ with $z_q \in \mathcal{B}_2$ for $1 \leq q \leq p$. Suppose that there exists $1 \leq q' \leq p$ such that $z_{q'} \notin \mathcal{B}^{(1)} = \{v^{11}(m, n) \mid m, n \in \mathbb{Z} \text{ with } m \leq n\}$. Let $q := \min\{1 \leq q' \leq p \mid z_{q'} \notin \mathcal{B}^{(1)}\}$. Then, z_q is either of the following form: $v^{1j}(m, n)$ for some $2 \leq j \leq d$ and $m \in \mathbb{Z}, n \in \mathbb{Z}_{\geq 0}$, or $v^{ij}(m, n) \in \mathcal{B}_+$ for some $2 \leq i \leq j \leq d$. Because $z_1, z_2, \dots, z_{q-1} \in \mathcal{B}^{(1)}$, it is obvious that $z_{q-1} \cdots z_1 u \in W \subset M_r^{(1)}$. Thus, using Lemma 4.6 below, we see that $z_q z_{q-1} \cdots z_1 u = 0$, which is a contradiction. Thus we conclude that $z_1, z_2, \dots, z_p \in \mathcal{B}^{(1)}$. Hence,

$$\begin{aligned} xu &= yzu = zu, & \text{since } y &= 1 \\ &= z_p z_{p-1} \cdots z_1 u \in W, & \text{since } z_1, z_2, \dots, z_p &\in \mathcal{B}^{(1)}. \end{aligned}$$

Since $xu \in \mathbb{C}\mathbf{1} \setminus \{0\}$, it follows that $\mathbf{1} \in W$, which implies that $W = M_r^{(1)}$. However, this is a contradiction, since W is assumed to be a proper $\mathcal{L}_r^{(1)}$ -submodule of $M_r^{(1)}$. Thus we have proved Lemma 4.5. $\square$

Let us show the following lemma, which has been used in the proof of Lemma 4.5.

Lemma 4.6. (1) Let $2 \leq j \leq d$, and $m \in \mathbb{Z}, n \in \mathbb{Z}_{\geq 0}$. Then, $v^{1j}(m, n)M_r^{(1)} = \{0\}$.
(2) Let $v^{ij}(m, n) \in \mathcal{B}_+$ with $2 \leq i \leq j \leq d$. Then, $v^{ij}(m, n)M_r^{(1)} = \{0\}$.

Proof. (1) Let $\mathcal{S}^{(1)}$ be the subset of $\mathcal{S}$ consisting of all finite sequences of elements in $\mathcal{B}_-^{(1)} := \mathcal{B}^{(1)} \cap \mathcal{B}_-$ that is weakly decreasing with respect to the total ordering $\succ$. Then, $\mathbb{B}^{(1)} := \{w(\mathbf{x}) \mid \mathbf{x} \in \mathcal{S}^{(1)}\} \subset \mathbb{B}$ is a linear basis of $M_r^{(1)}$. Therefore it suffices to show that

$v^{1j}(m, n)w(\mathbf{x}) = 0$ for all $\mathbf{x} \in \mathcal{S}^{(1)}$. Let $\mathbf{x} = (x_p \succeq x_{p-1} \succeq \cdots \succeq x_1) \in \mathcal{S}^{(1)}$ with $x_q \in \mathcal{B}^{(1)}$ for $1 \leq q \leq p$. We show $v^{1j}(m, n)w(\mathbf{x}) = 0$ by induction on the length p of the sequence $\mathbf{x}$. If $p = 0$, then the claim is obvious since $w(\mathbf{x}) = \mathbf{1}$. Assume that $p > 0$. Then,

$$\begin{aligned} v^{1j}(m, n)w(\mathbf{x}) &= v^{1j}(m, n)x_px_{p-1} \cdots x_1 \mathbf{1} \\ &= [v^{1j}(m, n), x_p]x_{p-1} \cdots x_1 \mathbf{1} + x_p \{v^{1j}(m, n)x_{p-1} \cdots x_1 \mathbf{1}\}. \end{aligned}$$

Since $v^{1j}(m, n)x_{p-1} \cdots x_1 \mathbf{1} = 0$ by the inductive assumption, the second term of the right-hand side is equal to 0. Assume that $x_p = v^{11}(s, t)$ with $s \leq t < 0$. By simple computation, we see that

$$\begin{aligned} [v^{1j}(m, n), x_p]_r &= [v^{1j}(m, n), v^{11}(s, t)]_r \\ &= \delta_{m+s, 0} m v^{1j}(t, n) + \delta_{m+t, 0} m v^{1j}(s, n). \end{aligned}$$

Hence it follows from the inductive assumption that $[v^{1j}(m, n), x_p]x_{p-1} \cdots x_1 \mathbf{1} = 0$. Thus we get $v^{1j}(m, n)w(\mathbf{x}) = 0$, thereby completing the proof of part (1).

(2) Since $2 \leq i \leq j \leq d$, it can be easily seen that $[v^{ij}(m, n), x] = 0$ for all $x \in \mathcal{B}^{(1)}$. Also, $v^{ij}(m, n)\mathbf{1} = 0$, since $v^{ij}(m, n) \in \mathcal{B}_+$. The assertion of part (2) follows immediately from these facts. This completes the proof of the lemma. $\square$

Proposition 4.1 follows from the results obtained in §§4.1 and 4.2. At the end of this section, we show the following proposition, which is needed in §6.3.

Proposition 4.7. *Assume that the $\mathcal{L}_r$ -module M_r is reducible, or equivalently, the $\mathcal{L}_r^{(1)}$ -module $M_r^{(1)}$ is reducible.*

- (1) *If W_1 is the maximal proper $\mathcal{L}_r^{(1)}$ -submodule of $M_r^{(1)}$, then $U(\mathcal{L}_r)W_1$ is the maximal proper $\mathcal{L}_r$ -submodule of M_r .*
- (2) *If W_2 is the maximal proper $\mathcal{L}_r$ -submodule of M_r , then $W_2 \cap M_r^{(1)}$ is the maximal proper $\mathcal{L}_r^{(1)}$ -submodule of $M_r^{(1)}$.*

Proof. We show that $U(\mathcal{L}_r)W_1 = W_2$ (for part (1)), and $W_1 = W_2 \cap M_r^{(1)}$ (for part (2)). Then we deduce from Lemma 4.5 that $U(\mathcal{L}_r)W_1$ is an $\mathcal{L}_r$ -submodule of M_r such that $U(\mathcal{L}_r)W_1 \neq M_r$. Hence we have $U(\mathcal{L}_r)W_1 \subset W_2$ by the maximality of W_2 . Also it follows from Lemma 4.2 (and the comment after it) that $W_2 \cap M_r^{(1)}$ is an $\mathcal{L}_r^{(1)}$ -submodule of $M_r^{(1)}$ such that $W_2 \cap M_r^{(1)} \neq M_r^{(1)}$. Hence we have $W_2 \cap M_r^{(1)} \subset W_1$ by the maximality of W_1 . Thus we obtain

$$U(\mathcal{L}_r)W_1 \cap M_r^{(1)} \subset W_2 \cap M_r^{(1)} \subset W_1.$$

Because it is obvious that $W_1 \subset U(\mathcal{L}_r)W_1 \cap M_r^{(1)}$, we get

$$U(\mathcal{L}_r)W_1 \cap M_r^{(1)} = W_2 \cap M_r^{(1)} = W_1,$$

which shows part (2).

Next, let us show $U(\mathcal{L}_r)W_1 = W_2$ (i.e., part (1)). Since $U(\mathcal{L}_r)W_1 \subset W_2$ as shown above, it suffices to show that $U(\mathcal{L}_r)W_1 \supset W_2$. Note that W_2 admits the weight space decomposition $W_2 = \bigoplus_{\lambda \in Q_+} (W_2)^\lambda$ with $(W_2)^\lambda = W_2 \cap (M_r)^\lambda$. Let $u \in W_2$ be a homogeneous element of weight $\lambda \in Q_+$, that is, $u \in (W_2)^\lambda$. We show by induction on $\theta(\lambda)$ that $u \in U(\mathcal{L}_r)W_1$ (for the definition of $\theta(\lambda)$, see (4.3)). If $\theta(\lambda) = 0$, then $u \in M_r^{(1)}$ by Remark 4.4, and hence $u \in W_2 \cap M_r^{(1)}$. Since $W_2 \cap M_r^{(1)} = W_1$ as shown above, it follows that $u \in W_1 \subset U(\mathcal{L}_r)W_1$. Next, let us assume that $\theta(\lambda) > 0$. Let $2 \leq i \leq d$ and $m \in \mathbb{Z}_{<0}$ be such that $\lambda(h_{i,m}) \geq 1$, and $N \in \mathbb{Z}_{<0}$ such that $\lambda(h_{1,N}) = 0$. Then we deduce from the proof of Lemma 4.2 that $v^{1i}(N, -m)u \neq 0$, and the weight λ' of $v^{1i}(N, -m)u$ satisfies $\theta(\lambda') = \theta(\lambda) - 1$. Hence it follows from the inductive assumption that $v^{1i}(N, -m)u \in U(\mathcal{L}_r)W_1$. Further, as in the proof of Lemma 4.2, we deduce that

$$v^{1i}(-N, m) \underbrace{v^{1i}(N, -m)u}_{\in U(\mathcal{L}_r)W_1} = mN \underbrace{\lambda(h_{i,m})}_{\geq 1} u,$$

which implies that $u \in U(\mathcal{L}_r)W_1$. This completes the proof of Proposition 4.7. $\square$

5 Irreducibility of $M_r^{(1)}$ for $r \in \mathbb{C} \setminus \mathbb{Z}$.

This subsection is devoted to proving the following proposition.

Proposition 5.1. *If $r \in \mathbb{C} \setminus \mathbb{Z}$, then $M_r^{(1)}$ is an irreducible $\mathcal{L}_r^{(1)}$ -module.*

5.1 Notation and some lemmas. For simplicity of notation, we set $v(m, n) := v^{11}(m, n)$ for $m, n \in \mathbb{Z}$, and $\Lambda_l := \Lambda_{1,l} \in \mathcal{H}^*$ for $l \in \mathbb{Z}_{<0}$. Recall that $\mathcal{S}^{(1)}$ denotes the subset of $\mathcal{S}$ consisting of all finite sequences of elements in $\mathcal{B}_-^{(1)} = \mathcal{B}^{(1)} \cap \mathcal{B}_-$ that is weakly decreasing with respect to the total ordering $\succ$, and $\mathbb{B}^{(1)} = \{w(\mathbf{x}) \mid \mathbf{x} \in \mathcal{S}^{(1)}\} \subset \mathbb{B}$ is a linear basis of $M_r^{(1)}$. For each $\lambda \in Q_+^{(1)}$, we denote by $\mathbb{B}_\lambda^{(1)}$ the set of all elements in $\mathbb{B}^{(1)}$ whose weight is equal to λ , and set $\mathcal{S}_\lambda^{(1)} := \{\mathbf{x} \in \mathcal{S}^{(1)} \mid w(\mathbf{x}) \in \mathbb{B}_\lambda^{(1)}\}$.

Let $u \in M_r^{(1)}$, and write it as a linear combination of elements of $\mathbb{B}^{(1)}$: $u = \sum_{b \in \mathbb{B}^{(1)}} \alpha_b b$ with $\alpha_b \in \mathbb{C}$ for $b \in \mathbb{B}^{(1)}$. Then we set $\mathbb{B}[u] := \{b \in \mathbb{B}^{(1)} \mid \alpha_b \neq 0\}$, and $\mathcal{S}[u] := \{\mathbf{x} \in \mathcal{S}^{(1)} \mid w(\mathbf{x}) \in \mathbb{B}[u]\}$.

The following formulas can be shown by simple computation.

Lemma 5.2. (1) *Let $s, t \in \mathbb{Z}_{>0}$, and $m, n \in \mathbb{Z}_{>0}$. Then,*

$$[v(-m, n), v(-s, -t)]_r = n\{\delta_{n,s}v(-m, -t) + \delta_{n,t}v(-s, -m)\}. \quad (5.1)$$

(2) *Let $s, t \in \mathbb{Z}_{>0}$ with $s \neq t$, and $m \in \mathbb{Z}_{>0}$. Then,*

$$[v(m, m), v(-s, -t)]_r = 2m\{\delta_{m,s}v(-t, m) + \delta_{m,t}v(-s, m)\}. \quad (5.2)$$

Lemma 5.3. *Let $m \in \mathbb{Z}_{>0}$, and $\nu \in \mathbb{Z}_{\geq 0}$. Then,*

$$v(m, m)v(-m, -m)^\nu \mathbf{1} = 2m^2\nu(r + 2\nu - 2)v(-m, -m)^{\nu-1} \mathbf{1}. \quad (5.3)$$

5.2 Proof of Proposition 5.1. We show that if $W \subset M_r^{(1)}$ is a nonzero $\mathcal{L}_r^{(1)}$ -submodule of $M_r^{(1)}$, then $W = M_r^{(1)}$. Suppose that $W \subsetneq M_r^{(1)}$. Because W admits the weight space decomposition $W = \bigoplus_{\lambda \in Q_+^{(1)}} W^\lambda$ with respect to $\mathcal{H}$ (see §4.2), it can be easily seen by usual way (see also Remark 4.3) that W contains a (homogeneous) singular vector u , i.e., a nonzero element u such that $u \in W^\lambda$ for some $\lambda \in Q_+^{(1)} \setminus \{0\}$, and $v(m, n)u = 0$ for all $m, n \in \mathbb{Z}$ with $m + n > 0$.

Claim 1. *The set $\{l \in \mathbb{Z}_{<0} \mid \lambda(h_l) > 0\}$ is identical to $\{-p, -p+1, \dots, -2, -1\}$ for some $p \in \mathbb{Z}_{\geq 1}$.*

Proof of Claim 1. Assume that $\{l \in \mathbb{Z}_{<0} \mid \lambda(h_l) > 0\} = \{l_p < l_{p-1} < \dots < l_1\}$; note that $l_q \leq -q$ for every $1 \leq q \leq p$. Suppose that $l_q < -q$ for some $1 \leq q \leq p$. We set $q_0 := \min\{1 \leq q \leq p \mid l_q < -q\}$. Since $\lambda(h_{l_{q_0}}) > 0$ and $\lambda(h_{-q_0}) = 0$, we deduce by a way similar to the proof of Lemma 4.2 that $v(l_{q_0}, q_0)u = 0$, and

$$\begin{aligned} v(l_{q_0}, q_0)v(-q_0, -l_{q_0})u &= l_{q_0}v(-q_0, q_0)u + q_0v(l_{q_0}, -l_{q_0})u \\ &= q_0 l_{q_0} \underbrace{\lambda(h_{-q_0})}_{=0} u + (-q_0 l_{q_0}) \underbrace{\lambda(h_{l_{q_0}})}_{>0} u \neq 0. \end{aligned}$$

Hence we get $v(-q_0, -l_{q_0})u \neq 0$. However, since $l_{q_0} < -q_0$, this contradicts the assumption that u is a singular vector. Thus we obtain $l_q = -q$ for all $1 \leq q \leq p$, thereby completing the proof of Claim 1.

Claim 2. *The weight λ of the singular vector u is of the form: $\lambda = \sum_{q=1}^p 2\nu_q \Lambda_{-q}$ with $\nu_q \in \mathbb{Z}_{>0}$ for $1 \leq q \leq p$. Furthermore, the set $\mathcal{S}[u]$ contains the element $\mathbf{x}_\lambda \in \mathcal{S}_\lambda^{(1)}$ such that*

$$w(\mathbf{x}_\lambda) = \prod_{1 \leq q \leq p} v(-q, -q)^{\nu_q} \mathbf{1} \in \mathbb{B}_\lambda^{(1)}.$$

Proof of Claim 2. For $\mathbf{x} \in \mathcal{S}[u]$ and $1 \leq q \leq p$, we define $\kappa_q(\mathbf{x})$ to be the number of $v(-q, -q)$ appearing in the sequence $\mathbf{x}$. Take $\mathbf{x} \in \mathcal{S}[u]$ such that the sum $\sum_{q=1}^p \kappa_q(\mathbf{x})$ is maximum, and assume that $w(\mathbf{x}) \in \mathbb{B}[u]$ is of the form:

$$w(\mathbf{x}) = \prod_{1 \leq q \leq p} v(-q, -q)^{\nu_q} \prod_{1 \leq t < s \leq p} v(-s, -t)^{\nu_{s,t}} \mathbf{1}$$

for some $\nu_q \in \mathbb{Z}_{\geq 0}$, $1 \leq q \leq p$, and $\nu_{s,t} \in \mathbb{Z}_{\geq 0}$, $1 \leq t < s \leq p$. Suppose that $\nu_{s_1, t_1} > 0$ for some $1 \leq t_1 < s_1 \leq p$. Then we prove that $v(-t_1, s_1)u \neq 0$. For this, it suffices to show

that the set $\mathbb{B}[v(-t_1, s_1)u]$ is not empty. Here we show that $\mathbb{B}[v(-t_1, s_1)u]$ contains the following element:

$$\prod_{1 \leq q \leq p, q \neq t_1} v(-q, -q)^{\nu_q} v(-t_1, -t_1)^{\nu_{t_1}+1} \prod_{\substack{1 \leq t < s \leq p, \\ (s,t) \neq (s_1, t_1)}} v(-s, -t)^{\nu_{s,t}} v(-s_1, -t_1)^{\nu_{s_1, t_1}-1} \mathbf{1}.$$

It can be easily seen by (5.1) that the element above is contained in $\mathbb{B}[v(-t_1, s_1)w(\mathbf{x})]$. On the other hand, we deduce, using (5.1) and the maximality of $\mathbf{x}$, that the element above is not contained in $\mathbb{B}[v(-t_1, s_1)w(\mathbf{y})]$ for all $\mathbf{y} \in \mathcal{S}[u]$ with $\mathbf{y} \neq \mathbf{x}$. Combining these, we conclude that the element above is contained in $\mathbb{B}[v(-t_1, s_1)u]$, and hence $v(-t_1, s_1)u \neq 0$. However, since $t_1 < s_1$, this contradicts the assumption that u is a singular vector. Thus we obtain $\nu_{s,t} = 0$ for all $1 \leq t < s \leq p$, thereby completing the proof of Claim 2.

By Claim 2, we may assume that the singular vector $u \in W^\lambda$ is of the form:

$$u = w(\mathbf{x}_\lambda) + \sum_{1 \leq t < s \leq p} \alpha_{s,t} w(\mathbf{x}_\lambda^{s,t}) + \sum_{\mathbf{x} \in \mathcal{S}'} \beta_{\mathbf{x}} w(\mathbf{x})$$

with $\alpha_{s,t} \in \mathbb{C}$ for $1 \leq t < s \leq p$, and $\beta_{\mathbf{x}} \in \mathbb{C}$ for $\mathbf{x} \in \mathcal{S}'$, where $\mathbf{x}_\lambda^{s,t}$ is the element of $\mathcal{S}_\lambda^{(1)}$ such that

$$w(\mathbf{x}_\lambda^{s,t}) = v(-s, -s)^{\nu_s-1} v(-t, -t)^{\nu_t-1} v(-s, -t)^2 \prod_{1 \leq q \leq p, q \neq t, s} v(-q, -q)^{\nu_q} \mathbf{1} \in \mathbb{B}_\lambda^{(1)},$$

and $\mathcal{S}' := \mathcal{S}_\lambda^{(1)} \setminus \{\mathbf{x}_\lambda, \mathbf{x}_\lambda^{s,t} \mid 1 \leq t < s \leq p\}$.

Claim 3. (1) We have $\alpha_{s,t} = -\nu_s$ for every $1 \leq t < s \leq p$.

(2) For every $1 \leq s \leq p$,

$$\nu_s(r + 2\nu_s - 2) + \sum_{1 \leq t \leq s-1} \alpha_{s,t} + \sum_{s+1 \leq t \leq p} \alpha_{t,s} = 0.$$

Proof of Claim 3. (1) Fix $1 \leq t < s \leq p$. We can easily check by using (5.1) that if the set $\mathbb{B}[v(-t, s)w(\mathbf{x})]$ with $\mathbf{x} \in \mathcal{S}_\lambda^{(1)}$ contains

$$w_1 := v(-s, -s)^{\nu_s-1} v(-s, -t) \prod_{1 \leq q \leq p, q \neq s} v(-q, -q)^{\nu_q} \mathbf{1},$$

then $\mathbf{x} = \mathbf{x}_\lambda$ or $\mathbf{x} = \mathbf{x}_\lambda^{s,t}$. By simple computation, along with (5.1), we get

$$v(-t, s)w(\mathbf{x}_\lambda) = 2\nu_s s w_1, \quad v(-t, s)w(\mathbf{x}_\lambda^{s,t}) = 2s w_1 + (\text{other term}).$$

Because $v(-t, s)w = 0$ by assumption, it follows that $2\nu_s s + 2s\alpha_{s,t} = 0$, and hence $\alpha_{s,t} = -\nu_s$.

(2) Fix $1 \leq s \leq p$. We can easily check by using (2.5), (5.1), and (5.2) that if the set $\mathbb{B}[v(s, s)w(\mathbf{x})]$ with $\mathbf{x} \in \mathcal{S}_\lambda^{(1)}$ contains

$$w_2 := v(-s, -s)^{\nu_s-1} \prod_{1 \leq q \leq p, q \neq s} v(-q, -q)^{\nu_q} \mathbf{1},$$

then $\mathbf{x} = \mathbf{x}_\lambda$, or $\mathbf{x} = \mathbf{x}_\lambda^{s,t}$ for $1 \leq t \leq s-1$, or $\mathbf{x} = \mathbf{x}_\lambda^{t,s}$ for $s+1 \leq t \leq p$. We see from Lemma 5.3 that

$$v(s, s)w(\mathbf{x}_\lambda) = 2s^2\nu_s(r + 2\nu_s - 2)w_2.$$

Also it follows from (5.2) that for each $1 \leq t \leq s-1$,

$$\begin{aligned} v(s, s)w(\mathbf{x}_\lambda^{s,t}) &= 2sX_1v(s, -t)v(-s, -t)v(-s, -s)^{\nu_s-1}\mathbf{1} \\ &\quad + 2sX_1v(-s, -t)v(s, -t)v(-s, -s)^{\nu_s-1}\mathbf{1} \\ &\quad + 2sX_1v(-s, -t)^2v(s, s)v(-s, -s)^{\nu_p-1}\mathbf{1}. \end{aligned} \quad (5.4)$$

Here, for simplicity of notation, we set

$$X_1 := \prod_{1 \leq q \leq p, q \neq t, s} v(-q, -q)^{\nu_q} v(-t, -t)^{\nu_s-1}.$$

The second and third terms of the right-hand side of (5.4) do not contribute the coefficient of w_2 since they contain $v(-s, -t)$. Also, the first term is:

$$\begin{aligned} 2sX_1v(s, -t)v(-s, -t)v(-s, -s)^{\nu_s-1}\mathbf{1} \\ = 2s^2X_1v(-t, -t)v(-s, -s)^{\nu_p-1}\mathbf{1} + (\text{other term}). \end{aligned}$$

Thus we obtain

$$v(s, s)w(\mathbf{x}_\lambda^{s,t}) = 2s^2w_2 + (\text{other terms})$$

for $1 \leq t \leq s-1$. Similarly, we can show that

$$v(s, s)w(\mathbf{x}_\lambda^{t,s}) = 2s^2w_2 + (\text{other terms})$$

for $s+1 \leq t \leq p$. Because $v(p, p)u = 0$ by assumption, we obtain

$$2s^2\nu_s(r + 2\nu_s - 2) + 2s^2 \sum_{1 \leq t \leq s-1} \alpha_{s,t} + 2s^2 \sum_{s+1 \leq t \leq p} \alpha_{t,s} = 0,$$

and hence the equation of part (2). Thus we have proved Claim 3.

Combining the equations in Claim 3 with $s = p$, we obtain $\nu_p(r + 2\nu_p - 2) - (p-1)\nu_p = 0$, and hence that

$$r + 2\nu_p - 1 - p = 0. \quad (5.5)$$

However, this is a contradiction, since r is assumed not to be an integer. This completes the proof of Proposition 5.1. $\square$

Remark 5.4. For later use, let us show the following assertion: Keep the notation in the proof of Proposition 5.1 above. Then, $\nu_1 = \nu_2 = \cdots = \nu_p$. Indeed, by the equations of Claim 3, we see that $\nu_s(r + 2\nu_s - 2) - (s - 1)\nu_s - \nu_{s+1} - \cdots - \nu_p = 0$ for every $1 \leq s \leq p$. Therefore,

$$\begin{aligned} & (\nu_s - \nu_{s+1})(r + 2\nu_{s+1} - s - 1 + 2\nu_s) \\ &= \left\{ \nu_s(r + 2\nu_s - 2) - (s - 1)\nu_s - \nu_{s+1} - \cdots - \nu_p \right\} \\ & \quad - \left\{ \nu_{s+1}(r + 2\nu_{s+1} - 2) - s\nu_{s+1} - \nu_{s+2} - \cdots - \nu_p \right\} \\ &= 0 \end{aligned}$$

for every $1 \leq s \leq p - 1$. Using this equation and (5.5), we can show by descending induction that $\nu_q = \nu_{q+1}$ for all $1 \leq q \leq p - 1$, and hence $\nu_1 = \nu_2 = \cdots = \nu_p$.

6 Reducibility of $M_r^{(1)}$ for $r \in \mathbb{Z}$.

6.1 Notation and proposition. For each $p \in \mathbb{Z}_{\geq 0}$, let $\mathbf{V}_p$ be the following matrix of size p with entries in $\mathcal{B}_-^{(1)}$:

$$\mathbf{V}_p = (v(-s, -t))_{1 \leq s, t \leq p}.$$

Since $xy = yx$ for all $x, y \in \mathcal{B}_-^{(1)}$, we can consider the determinant $\det \mathbf{V}_p$ of the matrix $\mathbf{V}_p$;

$$\det \mathbf{V}_p = \sum_{\sigma \in \mathfrak{S}_p} \text{sgn}(\sigma) \prod_{1 \leq q \leq p} v(-q, -\sigma(q)),$$

where $\mathfrak{S}_p$ denotes the symmetric group of degree p , and $\text{sgn}(\sigma)$ denotes the signature of a permutation $\sigma \in \mathfrak{S}_p$. In this subsection, we show the following proposition.

Proposition 6.1. *Assume that $r \in \mathbb{Z}$. Let $\nu \in \mathbb{Z}_{\geq 1}$ and $p \in \mathbb{Z}_{\geq 1}$ be positive integers satisfying the relation $r = 1 - 2\nu + p$. Then, $(\det \mathbf{V}_p)^\nu \mathbf{1}$ is a singular vector of $M_r^{(1)}$, that is, $v(m, n)(\det \mathbf{V}_p)^\nu \mathbf{1} = 0$ for all $m, n \in \mathbb{Z}$ with $m + n > 0$. Therefore the $\mathcal{L}_r^{(1)}$ -module $M_r^{(1)}$ is reducible.*

Theorem 2.6 follows immediately from Propositions 3.1, 3.4, 4.1, 5.1, and 6.1 (see also the comment after Theorem 2.6).

6.2 Proof of Proposition 6.1. Let us first show the following lemmas.

Lemma 6.2. *Let $m \in \mathbb{Z}$ with $1 \leq m \leq p$, and $n \in \mathbb{Z}_{\geq 0}$ with $m \neq n$. Then,*

$$[v(-m, n), \det \mathbf{V}_p] = 0. \tag{6.1}$$

Proof. If $n \geq p+1$ or $n = 0$, then the assertion is obvious, since $[v(-m, n), v(-s, -t)]_r = 0$ for all $1 \leq s, t \leq p$. Assume that $1 \leq n \leq p$. It can be seen from (5.1) that $[v(-m, n), v(-s, -t)]_r$ is equal to 0, or is contained in $\mathcal{B}_-^{(1)}$, up to a scalar multiple. Since $\text{ad } x$ is a derivation on $U(\mathcal{L}_r^{(1)})$ for $x \in \mathcal{L}_r^{(1)}$, we deduce that

$$[v(-m, n), \det \mathbf{V}_p] = \det \mathbf{W}_1 + \det \mathbf{W}_2 + \cdots + \det \mathbf{W}_p,$$

where $\mathbf{W}_s$, $1 \leq s \leq p$, is the matrix obtained by replacing the s -th row $(v(-s, -t))_{1 \leq t \leq p}$ of the matrix $\mathbf{V}_p$ with

$$\left([v(-m, n), v(-s, -t)]_r \right)_{1 \leq t \leq p}.$$

It follows from (5.1) that if $s \neq n$, then

$$\left([v(-m, n), v(-s, -t)]_r \right)_{1 \leq t \leq p} = (0, \dots, 0, nv(-s, -m), 0, \dots, 0),$$

where $nv(-s, -m)$ is placed at the n -th entry. Also, it follows from (5.1) that

$$\begin{aligned} & \left([v(-m, n), v(-n, -t)]_r \right)_{1 \leq t \leq p} - \underbrace{n (v(-m, -t))_{1 \leq t \leq p}}_{\text{the } m\text{-th row of } \mathbf{W}_n} \\ &= (0, \dots, 0, nv(-n, -m), 0, \dots, 0), \end{aligned}$$

where $nv(-n, -m)$ is placed at the n -th entry. Thus, $[v(-m, n), \det \mathbf{V}_p] = n \det \mathbf{V}'_p$, where $\mathbf{V}'_p$ is the matrix obtained by replacing the n -th column $(v(-s, -n))_{1 \leq s \leq p}$ of the matrix $\mathbf{V}_p$ with $(v(-s, -m))_{1 \leq s \leq p}$ (i.e., the m -th column of $\mathbf{V}_p$). Because the n -th and m -th columns of $\mathbf{V}'_p$ are equal, we get $\det \mathbf{V}'_p = 0$, thereby completing the proof of the lemma. $\square$

Lemma 6.3. *Let $1 \leq m \leq p$, and assume that $u \in M_r^{(1)}$ satisfies the conditions that $v(-t, m)u = 0$ for all $1 \leq t \leq p$ with $t \neq m$, and $v(-m, m)u = \alpha mu$ for some $\alpha \in \mathbb{C}$. Then we have*

$$v(m, m)(\det \mathbf{V}_p)u = 2m^2(2\alpha + r - p + 1) \det(\mathbf{V}_p^{(m)})u + (\det \mathbf{V}_p)v(m, m)u,$$

where $\mathbf{V}_p^{(m)}$ is the matrix obtained by removing the m -th row and the m -th column from the matrix $\mathbf{V}_p$.

Proof. We set $\mathfrak{S}_{p-1}^{(m)} := \{\sigma \in \mathfrak{S}_p \mid \sigma(m) = m\}$, which is a subgroup of $\mathfrak{S}_p$ isomorphic to the symmetric group $\mathfrak{S}_{p-1}$ of degree $p-1$. Let

$$R^{(m)} := \{1, (m, t) \mid 1 \leq t \leq p \text{ with } t \neq m\}$$

be a complete set of the representatives for the quotient $\mathfrak{S}_p/\mathfrak{S}_{p-1}^{(m)}$, where $(m, t) \in \mathfrak{S}_p$ denotes the transposition interchanging m and t . Decompose $\mathfrak{S}_p$ as: $\mathfrak{S}_p = \bigsqcup_{\tau \in \mathfrak{S}_{p-1}^{(m)}} \tau R^{(m)}$, with $\tau R^{(m)} := \{\tau, \tau \cdot (m, t) \mid 1 \leq t \leq p \text{ with } t \neq m\}$. Then we have

$$\begin{aligned} v(m, m)(\det \mathbf{V}_p) u &= [v(m, m), \det \mathbf{V}_p] u + (\det \mathbf{V}_p) v(m, m) u \\ &= \sum_{\tau \in \mathfrak{S}_{p-1}^{(m)}} \sum_{\sigma \in \tau R^{(m)}} \text{sgn}(\sigma) \left[v(m, m), \prod_{1 \leq s \leq p} v(-s, -\sigma(s)) \right] u + (\det \mathbf{V}_p) v(m, m) u. \end{aligned} \tag{6.2}$$

If $\tau \in \mathfrak{S}_{p-1}^{(m)}$, then we see, using (2.5), (5.2), and the assumption on u , that

$$\begin{aligned} &\left[v(m, m), \prod_{1 \leq s \leq p} v(-s, -\tau(s)) \right] u \\ &= \left\{ \prod_{1 \leq s \leq p, s \neq m} v(-s, -\tau(s)) \right\} [v(m, m), v(-m, -m)] u \\ &= \left\{ \prod_{1 \leq s \leq p, s \neq m} v(-s, -\tau(s)) \right\} \{4mv(-m, m) + 2rm^2\} u \\ &= 2m^2(2\alpha + r) \prod_{1 \leq s \leq p, s \neq m} v(-s, -\tau(s)) u. \end{aligned}$$

Assume that $\sigma = \tau \cdot (m, t) \in \tau R^{(m)}$ with $\tau \in \mathfrak{S}_{p-1}^{(m)}$ and $1 \leq t \leq p$ with $t \neq m$; note that $\sigma^{-1}(m) = t$, and $\sigma(s) = \tau(s)$ for $1 \leq s \leq p$ with $s \neq m, t$. Then,

$$\begin{aligned} &\left[v(m, m), \prod_{1 \leq s \leq p} v(-s, -\sigma(s)) \right] u \\ &= \left\{ \prod_{\substack{1 \leq s \leq p, \\ s \neq m, \sigma^{-1}(m)}} v(-s, -\sigma(s)) \right\} [v(m, m), v(-\sigma^{-1}(m), -m)v(-m, -\sigma(m))] u \\ &= \left\{ \prod_{1 \leq s \leq p, s \neq m, t} v(-s, -\tau(s)) \right\} [v(m, m), v(-t, -m)v(-m, -\tau(t))] u. \end{aligned}$$

Using (5.1), (5.2), and the assumption for u (note that $\tau(t) \neq m$ and $t \neq m$), we deduce that

$$\left[v(m, m), v(-t, -m)v(-m, -\tau(t)) \right] u = 2m^2 v(-t, -\tau(t)) u.$$

Thus,

$$\left[v(m, m), \prod_{1 \leq s \leq p} v(-s, -\sigma(s)) \right] u = 2m^2 \prod_{1 \leq s \leq p, s \neq m} v(-s, -\tau(s)) u.$$

Therefore, for each $\tau \in \mathfrak{S}_{p-1}^{(m)}$,

$$\begin{aligned} & \sum_{\sigma \in \tau R^{(m)}} \text{sgn}(\sigma) \left[v(m, m), \prod_{1 \leq s \leq p} v(-s, -\sigma(s)) \right] u \\ &= 2m^2 \left\{ (2\alpha + r) \text{sgn}(\tau) + \sum_{1 \leq t \leq p, t \neq m} \text{sgn}(\tau \cdot (1, t)) \right\} \prod_{1 \leq s \leq p, s \neq m} v(-s, -\tau(s)) u \\ &= 2m^2 (2\alpha + r - p + 1) \text{sgn}(\tau) \prod_{1 \leq s \leq p, s \neq m} v(-s, -\tau(s)) u. \end{aligned}$$

Combining this equation and equation (6.2), we see that

$$v(m, m)(\det \mathbf{V}_p) u = 2m^2 (2\alpha + r - p + 1) \det(\mathbf{V}_p^{(m)}) u + (\det \mathbf{V}_p) v(m, m) u,$$

as desired. $\square$

Proof of Proposition 6.1. It is obvious that $(\det \mathbf{V}_p)^\nu \mathbf{1}$ is a nonzero homogeneous element, and $(\det \mathbf{V}_p)^\nu \mathbf{1} \notin \mathbb{C} \mathbf{1}$. Let $m, n \in \mathbb{Z}$ be such that $m \leq n$ and $m + n > 0$; note that $n \geq 1$. If $n \geq p + 1$, then $v(m, n)(\det \mathbf{V}_p)^\nu \mathbf{1} = 0$ since the weight of $v(m, n)(\det \mathbf{V}_p)^\nu \mathbf{1} = 0$ is not contained in $Q_+^{(1)}$ (see (4.4)). Let us consider the case that $1 \leq n \leq p$.

Case 1. Assume that $1 \leq n \leq p$ and $-n < m < 0$. It follows from Lemma 6.2 that $[v(m, n), \det \mathbf{V}_p] = 0$. Hence

$$v(m, n)(\det \mathbf{V}_p)^\nu \mathbf{1} = (\det \mathbf{V}_p)^\nu v(m, n) \mathbf{1} = 0.$$

Case 2. Assume that $1 \leq n \leq p$ and $m = 0$. By direct computation, we see that $v(0, n) = (p + 1)^{-1} [v(n, p + 1), v(-p - 1, 0)]_r$. As mentioned above, we have $v(n, p + 1)(\det \mathbf{V}_p)^\nu \mathbf{1} = 0$. Also, since $[v(-s, -t), v(-p - 1, 0)]_r = 0$ for all $1 \leq s, t \leq p$, it follows that $[v(-p - 1, 0), \det \mathbf{V}_p] = 0$, and hence that $v(-p - 1, 0)(\det \mathbf{V}_p)^\nu \mathbf{1} = 0$ as above. Thus we get $v(0, n)(\det \mathbf{V}_p)^\nu \mathbf{1} = 0$.

Case 3. Assume that $1 \leq n \leq p$ and $1 \leq m \leq n$. First let us consider the case that $m = n$. Namely, we show that $v(m, m)(\det \mathbf{V}_p)^\nu \mathbf{1} = 0$ for all $1 \leq m \leq p$. It follows

immediately from Lemma 6.2 that $u = (\det \mathbf{V}_p)^{\nu_1} \mathbf{1}$, $\nu_1 \in \mathbb{Z}_{\geq 0}$, satisfies the assumption of Lemma 6.3 with $\alpha = 2\nu_1$. Hence, by using Lemma 6.3 repeatedly, we obtain

$$\begin{aligned}
v(m, m)(\det \mathbf{V}_p)^{\nu} \mathbf{1} &= v(m, m)(\det \mathbf{V}_p)(\det \mathbf{V}_p)^{\nu-1} \mathbf{1} \\
&= 2m^2 \{4(\nu-1) + r - p + 1\} (\det \mathbf{V}_p^{(m)})(\det \mathbf{V}_p)^{\nu-1} \mathbf{1} + (\det \mathbf{V}_p) v(m, m)(\det \mathbf{V}_p)^{\nu-1} \mathbf{1} \\
&= \dots\dots\dots \\
&= 2m^2 (\det \mathbf{V}_p^{(m)}) \left\{ \sum_{\nu_1=0}^{\nu-1} (4\nu_1 + r - p + 1) \right\} (\det \mathbf{V}_p)^{\nu-1} \mathbf{1} \\
&= 4m^2 \nu (2\nu - 1 + r - p) (\det \mathbf{V}_p^{(m)})(\det \mathbf{V}_p)^{\nu-1} \mathbf{1}.
\end{aligned}$$

Since $2\nu - 1 + r - p = 0$ by assumption, we get $v(m, m)(\det \mathbf{V}_p)^{\nu} \mathbf{1} = 0$.

Next, let us consider the case that $m < n$. By direct computation, we see that $v(m, n) = (2m)^{-1} [v(m, m), v(-m, n)]_r$. Since $v(-m, n)(\det \mathbf{V}_p)^{\nu} \mathbf{1} = 0$ by Case 1, and since $v(m, m)(\det \mathbf{V}_p)^{\nu} \mathbf{1} = 0$ by the argument above, it follows that $v(m, n)(\det \mathbf{V}_p)^{\nu} \mathbf{1} = 0$. This completes the proof of Proposition 6.1 $\square$

6.3 Irreducibility of the quotient modules. Fix $r \in \mathbb{Z}$ as above. Denote by $W_r^{(1)}$ the $\mathcal{L}_r^{(1)}$ -submodule of $M_r^{(1)}$ generated by the singular vectors obtained in Proposition 6.1, i.e.,

$$W_r^{(1)} := \left\langle (\det \mathbf{V}_p)^{\nu} \mathbf{1} \mid p, \nu \in \mathbb{Z}_{\geq 1} \text{ with } r = 1 - 2\nu + p \right\rangle \subset M_r^{(1)}.$$

Proposition 6.4. *A singular vector of $M_r^{(1)}$ is equal to a scalar multiple of the singular vector $(\det \mathbf{V}_p)^{\nu} \mathbf{1}$ for some $p, \nu \in \mathbb{Z}_{\geq 1}$ with $r = 1 - 2\nu + p$. Therefore the $\mathcal{L}_r^{(1)}$ -submodule $W_r^{(1)}$ is the maximal proper submodule of $M_r^{(1)}$, and hence the quotient $\mathcal{L}_r^{(1)}$ -module $M_r^{(1)}/W_r^{(1)}$ is irreducible.*

Proof. Let $u \in M_r^{(1)}$ be a singular vector, and assume that the weight of u is equal to $\lambda \in \mathcal{Q}_+^{(1)} \setminus \{0\}$. By Claim 2 in the proof of Proposition 5.1 and by Remark 5.4, we see that the weight λ is of the form: $2\nu \sum_{q=1}^p \Lambda_{-q}$ for some $p > 0$ and $\nu > 0$, and that $w(\mathbf{x}_{\lambda}) \in \mathbb{B}[u]$. In addition, it follows from (5.5) that $r = 2\nu - 1 - p$. Thus, by Proposition 6.1, $(\det \mathbf{V}_p)^{\nu} \mathbf{1}$ is a singular vector. Because $w(\mathbf{x}_{\lambda}) \in \mathbb{B}[(\det \mathbf{V}_p)^{\nu} \mathbf{1}]$ by definition, there exists $\alpha \in \mathbb{C} \setminus \{0\}$ such that

$$w(\mathbf{x}_{\lambda}) \notin \mathbb{B}[u - \alpha(\det \mathbf{V}_p)^{\nu} \mathbf{1}]. \quad (6.3)$$

Here we should remark that $u - \alpha(\det \mathbf{V}_p)^{\nu} \mathbf{1}$ is also a singular vector of weight λ if it is nonzero. Therefore we deduce from (6.3) and Claim 2 in the proof of Proposition 5.1 that $u - \alpha(\det \mathbf{V}_p)^{\nu} \mathbf{1} = 0$, and hence $u = \alpha(\det \mathbf{V}_p)^{\nu} \mathbf{1}$. Thus we have proved the first assertion of the proposition. The other assertions are obvious. This completes the proof of the proposition. $\square$

Let I_r be the ideal of the VOA $V_{\mathcal{J}}$ ($= M_r$) generated by all $(\det \mathbf{V}_p)^\nu \mathbf{1}$ for $p, \nu \in \mathbb{Z}_{\geq 1}$ with $r = 1 - 2\nu + p$.

Corollary 6.5. *The ideal I_r is the maximal proper ideal of the VOA $V_{\mathcal{J}}$. Therefore the quotient VOA $V_{\mathcal{J}}/I_r$ is simple.*

Proof. It follows from Propositions 4.7 and 6.4 that $W_r := U(\mathcal{L}_r)W_r^{(1)}$ is the maximal proper $\mathcal{L}_r$ -module of M_r . Then we see by Corollary 3.6 that W_r is the maximal proper ideal of the VOA $V_{\mathcal{J}}$. So, let us show that $I_r = W_r$. The inclusion $I_r \subset W_r$ follows from the fact that the ideal W_r contains all $(\det \mathbf{V}_p)^\nu \mathbf{1}$ for $p, \nu \in \mathbb{Z}_{\geq 1}$ with $r = 1 - 2\nu + p$. Thus, I_r is a proper ideal of $V_{\mathcal{J}}$, which implies that I_r is a proper $\mathcal{L}_r$ -submodule of M_r (see the proof of Proposition 3.4). Since I_r contains all $(\det \mathbf{V}_p)^\nu \mathbf{1}$ for $p, \nu \in \mathbb{Z}_{\geq 1}$ with $r = 1 - 2\nu + p$, it can be easily seen from the definition of W_r that $I_r \supset W_r$. Thus we get $I_r = W_r$, thereby completing the proof of the corollary. $\square$

References

- [Al1] A.A. Albert, On Jordan algebras of linear transformations, *Trans. Amer. Math. Soc.* **59** (1946), 524–555.
- [Al2] A.A. Albert, A structure theory for Jordan algebras, *Ann. of Math.* (2) **48** (1947), 546–567.
- [As] T. Ashihara, On vertex operator algebras whose Griess algebras are Jordan algebras, doctor thesis, 2008.
- [AM] T. Ashihara and M. Miyamoto, Deformation of central charges, vertex operator algebras whose Griess algebras are Jordan algebras, arXiv:0806.4308, to appear in *J. Algebra*.
- [B] R.E. Borcherds, Vertex algebras, Kac-Moody algebras, and the Monster, *Proc. Nat. Acad. Sci. U.S.A.* **83** (1986), 3068–3071.
- [DMZ] C. Dong, G. Mason, and Y. Zhu, Discrete series of the Virasoro algebra and the moonshine module, in “Algebraic Groups and their Generalizations: Quantum and Infinite-dimensional Methods” (W.J. Haboush and B.J. Parshall, Eds.), Proc. Sympos. Pure Math. Vol. 56 Part 2, pp. 295–316, Amer. Math. Soc., Providence, RI, 1994.
- [FK] J. Faraut, and A. Korányi, “Analysis on Symmetric Cones”, Oxford Mathematical Monographs, Oxford University Press, New York, 1994.
- [FLM] I.B. Frenkel, J. Lepowsky, and A. Meurman, “Vertex Operator Algebras and the Monster”, Pure and Appl. Math. Vol. 134, Academic Press, 1988.
- [FZ] I.B. Frenkel and Y. Zhu, Vertex operator algebras associated to representations of affine and Virasoro algebras, *Duke Math. J.* **66** (1992), 123–168.
- [G] R.L. Griess, The friendly giant, *Invent. Math.* **69** (1982), 1–102.

- [J] N. Jacobson, “Structure and representations of Jordan algebras”, American Mathematical Society Colloquium Publications Vol. XXXIX, American Mathematical Society, Providence, R.I., 1968.
- [L1] C.H. Lam, Construction of vertex operator algebras from commutative associative algebras, *Comm. Algebra* **24** (1996), 4339–4360.
- [L2] C.H. Lam, On VOA associated with special Jordan algebra, *Comm. Algebra* **27** (1999), 1665–1681.
- [LL] J. Lepowsky and H. Li, “Introduction to Vertex Operator Algebras and Their Representations”, Progr. Math. Vol. 227, Birkhäuser Boston, Boston, 2004.
- [M] M. Miyamoto, Griess algebras and conformal vectors in vertex operator algebras, *J. Algebra* **179** (1996), 523–548.
- [W] W. Wang, Rationality of Virasoro vertex operator algebras, *Int. Math. Res. Not.* **1993** (1993), no. 7, 197–211.